

WEB DIRECTION

7.85"(h) x 4.3"(w)



INSECT KILLER

CONCENTRATE

PEOPLE & PETS MAY RE-ENTER

BOUND

BOOKLET

TREATED AREA
AFTER SPRAY DRIES

GUARANTEED RESULTS OR YOUR MONEY BACK

TO USE	Lawns, Trees and Shrubs, Roses and Flowers, Up to 2 feet of outside foundation surface of buildings
KILLS	Ants, Mosquitoes, Gypsy moth, bagworm, budworm & leafroller, armyworms cutworms, fall webworm, sod webworm, tent caterpillar, leafminer, Millipedes, Springalias, Silverfish, firebrats, Earnigs, Cockroaches, Fleas, Spiders, Grasshoppers, Crickets, Chinch bug, boxelder bug, plant bug, lace bug, Leafhoppers, spittlebugs, psyllids, whitefly, aphids, cicadas, scales, mealybug, adelgids, treehoppers, Flies, Blacklegged and Brown dog ticks, Spotted Lanternify' and Other Listed insects.
WHEN TO USE	To treat lawns and vegetation: When insects or damage first appear. When using as perimeter treatment: Early in season or when insects appear
HOW MUCH TO USE	For Lawns and perimeter treatments: Dilute 4 fl oz of this product per gallon o water for 0.011% solution to evenly cover 1000 sq ft of area (0.000938 lbs ai per 1000 sq ft). For Ornamentals, Trees and Flowers: Dilute 4 fl oz of this product per gallon of water for 0.011% solution and fully spray foliage (0.000938 lbs ai per 1000 sq ft). For Fleas, Mosquitoes, Scorpions, Mole Crickets, Spiders, Crickets & Grasshoppers and Blacklegged and Brown dog ticks: Dilute 15 fl oz of this product per gallon of water for 0.04% solution to evenly cover 1,000 sq ft of area (0.003516 lbs ai per 1000 sq ft).
COVERAGE	MAKES UP TO TREATS UP TO 32,000 SQ. SQ.

*Not Registered for Use by California

Distributed by: Central Garden & Pet, Garden Division 1000 Parkwood Circle, Suite 700 Atlanta. GA 30339 EPA Reg. No. 279-3347-90098 EPA Est. No. 92162-MO-001,, 33595-MO-6₈, 67572-GA-1₁, 67572-GA-2₂, 59144-GA-1₅ Subscript used is first letter of Run Code on container.

 $See\ Inside\ Back\ Booklet\ for\ Precautionary\ Statements,\ First\ Aid\ and\ Complete\ Directions\ For\ Use.$

300536557 | AM1242 v7 - 1/25



AMDRO.com

1-800-265-0761 | gardendivision@central.com

CENTRAL GARDEN & PET

Amdro Quick Kill, Central Garden & Pet and Central Garden & Pet with design are trademarks of Central Garden & Pet Company.

WEB DIRECTION

7.85"(h) x 4.3"(w)



CONCENTRATE

FIRST AID

If swallowed

- Call poison control center or doctor immediately for treatment advice.
 Have person sip a glass of water if able to swallow.
 - DO NOT induce vomiting unless told to do so by the poison control
 - DO NOT give anything by mouth to an unconscious person.

If in eyes

7.59"

- Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then
- continue rinsing eve. Call a poison control center or doctor for treatment advice.

HOTLINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1- (800)-265-0761 for Emergency Assistance.

NOTE TO PHYSICIAN

Skin contact (vapor or powder) may be followed by transient tingling or numbness, usually of the face, but this subsides without treatment. This product is a pyrethroid. If large amounts have been ingested, the stomach and intestines should be evacuated. Treatment is

For Information Regarding the Use of this Product Call 1-800-265-0761.

PRECAUTIONARY STATEMENTS

ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to fish, aquatic invertebrates, oysters and shrimp. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems. Care should be used when spraying to avoid fish and reptile pets in/around ornamental ponds. This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are foraging the

STORAGE AND DISPOSAL

Pesticide Storage: Store in original container in a cool dry area away from heat or open flame and in an area inaccessible to children and pets. Keep from freezing.

DO NOT pour or dispose down-the-drain or sewer. Call your local solid waste agency for local disposal options.

PESTICIDE DISPOSAL AND CONTAINER HANDLING: Nonrefillable Plastic container. Do not reuse or refill this container. If empty: Place in trash or offer for recycling if available. If partly filled: Call your local solid waste agency for disposal instructions. Never place unuséd product down any indoor or outdoor drain.

Distributed by: Central Garden & Pet, Garden Division 1000 Parkwood Circle, Suite 700 Atlanta, GA 30339

EPA Reg. No. 279-3347-90098 EPA Est. No. 92162-MO-001, 33595-MO-6, 67572-GA-1,, 67572-GA-2,, 59144-GA-1, Subscript used is first letter of Run Code on container



AMDR().com

1-800-265-0761 | gardendivision@central.com

- Amdro Quick Kill, Central Garden & Pet and
- Central Garden & Pet with design are trademarks of Central Garden & Pet Company. 300536557 | AM1242 v7 1/25

- 3**.**625" -

BASE TEXT (Underneath Book)

BASE LABEL

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. SHAKE WELL BEFORE USINĞ

USE RESTRICTIONS

- For outdoor use only.
- DO NOT treat pets with this product.
- DO NOT apply this product in a way that will contact adults, children, or pets, either directly or through drift.
- DO NOT allow adults, children, or pets to enter the treated area until sprays
- Remove pets, birds, and cover fish aquariums before spraying.
- Avoid contamination of food and feedstuffs. Remove or cover exposed food and drinking water before application.
- DO NOT use in food handling establishments in which food is held, processed, prepared, and/or served.
- DO NOT apply directly to animals.
- DONOT use in aircraft.

7.1383"

FOR OUTDOOR USES

All outdoor applications must be limited to spot or crack-and-crevice treatments only, except for the following permitted uses:

- 1. Application to pervious surfaces such as soil, lawn, turf, and other vegetation:
- 2. Perimeter band treatments of 7 feet wide or less from the base of a man-made structure to pervious surfaces (e.g., soil, mulch, or lawn);
- 3. Applications to underside of eaves, soffits, doors, or windows permanently protected from rainfall by a covering, overhang, awning, or other structure
- Applications around potential exterior pest entry points into man-made structures such as doorways and windows, when limited to a band not to
- 5. Applications to vertical surfaces (such as the side of a man-made structure) directly above impervious surfaces (e.g., driveways, sidewalks, etc.), up to
- 6. Applications to vertical surfaces directly above pervious surfaces, such as soil, lawn, turf, mulch or other vegetation only if the pervious surface does not drain into ditches, storm drains, gutters, or surface waters.

For spot treatments: Spot treatments must not exceed two square feet in size (for example, 2 ft by 1 ft or 4 ft by 0.5 ft).

For crack-and-crevice treatments: To treat insects harbored in voids and cracks-and-crevices, applications must be made in such a manner to limit dripping and avoid runoff onto untreated structural surfaces and plants. DO NOT apply directly to impervious horizontal surfaces such as sidewalks, driveways, and patios except as a spot or crack-and-crevice treatment. Treat surfaces to ensure thorough coverage but avoid runoff.

DO NOT make applications during rain. Avoid making applications when rainfall is expected before the product has sufficient time to dry (minimum 4 hours). Rainfall within 24 hours after application may cause unintended runoff of pesticide application.

DO NOT apply or irrigate to the point of runoff.
DO NOT apply the product into fish pools, ponds, streams, or lakes. DO NOT apply directly to sewers or storm drains, or to any area like a drain or gutter where drainage to sewers, storm drains, water bodies, or aquatic habitat can occur. For soil or foliar applications, DO NOT apply by ground within 25 feet of lakes, reservoirs, rivers, permanent streams, marshes or natural ponds, estuaries and commercial fish farm ponds.

DO NOT allow the product to enter any drain during or after application.

DO NOT allow to enter indoor or outdoor drains. No permita la entrada a desagües internos o externos

NOTE: The application of this product to vinyl and aluminum siding (particularly lightly colored, aged, weathered or otherwise damaged) may result in staining, bleaching or discoloration. Factors including extreme heat and direct sunlight can promote damage. Before applying this product to vinyl or aluminum siding, treat a small area and evaluate 30 minutes later to allow any potential staining to occur. Regardless of the test results, do not apply to vinyl or aluminum siding while exposed to direct sunlight or during the heat of the day.

MIXING INSTRUCTIONS - HOW MUCH TO USE

This product is to be diluted with water for spray application.

- 1) Fill sprayer with the desired volume of water and add Amdro Quick Kill® Insect Killer For Lawn & Landscape Concentrate.
- 2) To prepare a 0.003% ready to use solution, dilute 1fl oz of this product per gallon of water.
- To prepare a 0.011% ready to use solution, dilute 4fl oz of this product per gallon of water.
- To prepare a 0.02% ready to use solution, dilute 7.5 fl oz of this product per gallon of water.
- To prepare a 0.04% ready to use solution, dilute 15 fl oz of this product per gallon of water.

3) Mix only the amount of solution needed for the application.

4) Close and shake before use in order to insure proper mixing

HOW TO USE - OUTDOORS

NUISANCE PESTS ON OUTDOOR SURFACES

To Kill Ants, Earwigs, and other named Pests on Outside Surfaces and Around Buildings (except for Fleas, Mosquitoes, Spiders, Scorpions, Crickets, Mole Crickets, and Grasshoppers - see specific instructions below)

- Apply 0.003% solution to point of run-off as a residual spray to vegetation including bushes and trees around buildings and other structures.
- For ants, apply to ant trails, around doors and windows and other places where ants may be found.
- To prevent pests from entering buildings, apply 0.003% solution to a band of soil and vegetation up to 7 feet wide around and adjacent to the building. Also, treat the building foundation to a height of up to 2 feet where pests are active and may find entrance. Apply as a coarse spray. Thoroughly and uniformly wet the band area, using 1 gallon of spray solution per 1,000 sq ft (0.000234 lb ai/1000 sq ft).
- DO NOT apply if surface is not dry at the time of application. Apply to attics and unoccupied lofts at the same time and at the same rate.

To Kill Fleas, Mosquitoes, Spiders, Scorpions, Mole Crickets, Crickets and Grasshoppers

- · For best results, locate areas where these pests are most active or may become
- Treat using a 0.04% solution (15 fl oz per gallon of water per 1,000 sq ft)
- (0.003516 lbs ai/1,000 sq ft). Use a coarse spray to ensure thorough coverage. Around buildings, apply an uninterrupted band up to 7 feet wide around the perimeter of the building and up to 2 feet along the side or foundation of
- Use a spot and crack and crevice treatment around potential entry points including doors, windows, and eaves.

To Kill Bees and Wasps Outside

- Apply to nests in the evening when insects are at rest.
- Thoroughly spray nest and entrance and surrounding areas where insects alight.

Summary Table for Wood-Infesting Insects:	
Where to Use	Outdoor and home perimeter
Kills	Carpenter ants, Crazy Ants, Tawny Crazy Ants, termites, Blacklegged and Brown dog ticks, Carpenter bees, wood boring beetles and other listed wood-destroying pests
When to Use	To kill wood-destroying insects outside, including home perimeters, decks, wood piles, wood patios, wood sheds and other wooded structures. Also, kills Blacklegged and Brown dog ticks on outdoor surfaces: When insects first appear

To Kill Carpenter Ants Outdoors

 Apply to carpenter ant trails around doors and windows and other places where carpenter ants have been observed or are expected to forage.

For best results, locate and treat carpenter ant nests.

To Kill Carpenter Ants Outdoors (cont'd)

Apply a perimeter treatment as described above.

To protect firewood from carpenter ants, apply to the soil beneath where the firewood will be stacked. Use 1 gallon of 0.04% spray solution to treat 1000 sq ft (0.003516 lbs ai/1000 sq ft). DO NOT treat firewood with this product.

To Kill Termites Outdoors

To protect firewood from termites, apply to the soil beneath where the firewood will be stacked. Use 1 gallon of 0.04% spray solution (15 fl oz per gallon water) to treat 1000 sq ft (0.003516 lbs ai/1000 sq ft). DO NOT treat firewood with this product.

DO NOT use this product as the sole source of control for active, structural infestations by subterranean termites; the purpose of this application is to kill workers or winged reproductive forms of termites which are present at the time of treatment. It is not intended to provide structural pest control. It is not a substitute for mechanical alteration, soil and foundation treatment, but merely a supplement. For active, structural infestations by subterranean termites, this product can only be used to supplement a federally registered conventional product that is registered as a sole source for termite control. This product will not eliminate termite infestations or provide protection against future infestation.

To Kill Blacklegged and Brown Dog Ticks Outdoors • Treat using a 0.04% solution (15 fl oz per gallon of water per 1,000 sq ft) (0.003516 lbs ai/1,000 sq ft). Use a coarse spray to ensure thorough coverage.

To prevent infestation of the home, treat soil and vegetation around the exterior perimeter of the home as described above.

Treat soil and vegetation adjacent to wooded areas on the property where blacklegged and brown dog ticks may be harboring.

Treat areas of taller vegetation up to 2 ft, around the exterior of the property.

Spray to wet, do not apply to the point of runoff.

Apply thoroughly and uniformly.

Treatment of Wood-in-Place to Kill Exposed Stages of Wood-Infesting Insects

For the control of insects including Termites, Carpenter Bees, and Carpenter Ants in localized areas of infested wood in and around structures, apply to voids and galleries in damaged wood and in spaces between wooden members of a structure and between wood and foundations where wood is vulnerable. Not intended to replace structural treatments.

LAWNS

7.1383"

To Kill Ants, Fire Ants, Beetles, Chinch Bugs, Sod Webworms, Weevils and other listed Pests in Lawns

Uniformly apply one gallon of 0.02% solution over a 1,000 sq ft area (0.001758 lbs ai/1000 sqft) with a sprayer

If possible, mow before application. DO NOT apply if lawn is longer than 3 inches at the time of application.

Repeat application is necessary only if there are signs of renewed insect activity.

To Kill Fleas and Mosquitoes

For best results, locate areas where these pests are most active or may become

Treat using 15 fl. oz. per gallon of water per 1,000 sq ft (0.003516 lbs ai/1000 sq ft). Apply uniformly over lawn. Treat area directly using a coarse spray until thoroughly wet. Do not apply to point of runoff.

Ant Mound Treatments

To treat ant mounds including imported fire ant mounds, dilute 0.8 fl oz of this product in 1-2 gal water and apply to each mound area by sprinkling the mound until it is wet and treat a 4foot diameter circle around the mound. Use the higher volume for mounds larger than 12". For best results, apply in cool weather, including in early morning or late evening hours, but not in the

ORNAMENTALS. ROSES. FLOWERS & TREES

To Kill Ants, (including cut ants), Aphids, Bagworms, Earwigs, Sow Bugs, Millipedes, Beetles, Weevils, Plant Bugs, Leaf-cutting Bees, Sawflies, Caterpillars, spotted lanternfly', Moths and other listed pests on Ornamentals, Roses, Flowers and Trees (including deciduous and evergreen trees, shrubs, ground covers, bedding plants, annuals, perennials, and foliage plants).

Apply 0.02-0.04% solution to point of run-off as a spray. Spray up to the point of run-off making sure to cover all sides of leaves and trunk. Retreat if pests return.

For ground covers, bedding plants, foliage, annuals and perennials, apply one gallon of $0.02 \cdot 0.04\%$ solution per 1,000 sq ft as a uniform spray (0.001758 - 0.003516 lbs ai/sq ft). Repeat treatment as necessary to achieve control using higher labeled rates as pest pressure & foliage area increases. Do not repeat application no more than once per seven days. NOTE: Certain plants may be sensitive to the final spray solution. Treat a limited area and observe for one week prior to application to the entire planting.

Apply directly to bark and foliage where insect is sighted.
 Fully spray foliage, stems and bark until is wet.

PESTS CONTROLLED

Caterpillars, Butterflies and Moths

Tent forming caterpillars, cocoon forming caterpillars, chrysalis forming

Leaf feeding caterpillars (including leaves of flowering plants, bushes and Lear reeding caterpillars (including leaves of nowering plants, busnes and other ornamentals, vegetables and trees), azalea leafminers, green cloverworm, fruit eating caterpillars, legume feeding caterpillars, vegetable feeding caterpillars, and leafrolling caterpillars. Caterpillars of the Yucca Moth family including Fairy moths and Yucca moths; Clothes moth family, bagworm moth family, Gelechiid moth family including Potato tuber moths; Ermine Moth family including Potato tuber moths; Ermine Moth family including Managered. Ailanthus Webworm moth; Clear Winged moth family including Manroot borer, Squash Vine borer, Doll's Clearwing moth, California Sycamore borer; Tortricid Moth family (millers) including Fruit Tree Leafroller moth, Orange Tortrix, and Codling moth, Skippers including Least Skipper, Silver-spotted Skipper, and Long-tail Skipper; Swallowtails including Green swallowtail, Zebra Swallowtail, Giant Swallowtail (Orange Tail, Orange Puppy), Pale Swallowtail, Eastern Tiger Swallowtail, Eastern Black Swallowtail, Spicebush Swallowtail, and Phoebus; Whites, Sulphurs and Orange Tips family including Sara Orange Tip, Great Southern White, and European Cabbage Butterfly; Gossamer-Winged Butterflies Family including Great Purple Hairstreak, Spring Azure, Early Hairstreak, Eastern Tailed blue, Harvester, Silvery Blue, Brown Elfin, Marine Blue, American Copper, and Gray Hairstreak; Metalmark family including Northern Metalmark; Snout Butterflies family including Common Snout Butterfly; Brush-Footed Butterfly family including Gulf Fritillary, White Peacock, Harris' Butterrily ramily including bull Fritiliary, White Peacock, Harris
Checkerspot, Painted Lady (Thistle butterfly), American Painted Lady
(Painted Beauty), Baltimore, Zebra (Yellow-barred Heliconian), Viceroy,
White Admiral (Banded Purple), Mourning Cloak, Milbert's Tortoise Shell,
Pearl Crescent, Question Mark, Buckeye, Great Spangled Fritillary, and Red
Admiral; Hackberry and Goatweed Butterfly family including Hackberry
butterfly; Satyrs, Nymphs and Arctics family including Wood Nymph
(American Grayling), White Wood Satyr, Eyed Brown (Grass Nymph), Pearly Eye, and Chryxus Arctic; Smoky Moth family including Grape Leaf Skeletonizer; Slug Caterpillar Moth family including Saddleback Caterpillar Moth; Pyralid Moths family including Mediterranean Flour Moth Sod Webworm Moth, Indian Meal Moth, Meal Moth; Plume Moth family including Artichoke Plume Moth; Measuringworm Moth family including Tulip Tree Beauty, California Cankerworm Moth, and Large California Spanworm Moth; Tent Caterpillar Moth family including Western Tent Caterpillar Moth; Giant Silkworm Moth family including Luna Moth, Polyphemus moth, Io Moth, Promethea Moth (Spicebush Silkmoth), Regal

Caterpillars, Butterflies and Moths (cont'd)

Moth (Royal Walnut Moth), Rosy Maple Moth, Imperial Moth, Sheep Moth, Cecropia Moth, and Cynthia Moth; Sphinx moth Sneep Moth, Lecropia Moth, and Lynthia Moth; Spfinix Moth family including Virginia-creeper Sphinx, Pandora Sphinx, Hummingbird Moth (Common Clearwing), White-lined Sphinx, Tomato Hornworm Moth, Tobacco Hornworm Moth, Big Poplar Sphinx, Cerisy's Sphinx, and Wild-cherry Sphinx; Prominents family including Tentacled Prominenets (Puss Moths), and Red-humped Appleworm Moth; Tiger Moth family including Ornate Tiger Moth, Yellow Woolly Bear Moth, Acraea Moth, Mikwed Tiger Moth, Yellow Woolly Bear Moth, Acraea Moth, Mikwed Tiger Moth, Pontad Tiger Moth Calear Weelly Report Cataroliles Moth (Pandod Spotted Tiger Moth, Colona, Woolly Bear Caterpillar Moth (Banded Woolly Bear), Lichen Moth, and Rattlebox Moth; Ctenuchid Moth family including Virginia Ctenuchid moth; Tussok moth family including Gypsy Moth, White-marked Tussok Moth; Owlet Moth family including Eight-spotted Forester, Black Witch, Alfalfa Looper, Sweetheart Underwing, Locust Underwing, and Hebrew.

Common Bristletails

Silverfish Family including Silverfish, and Firebrat

Podurid Springtail family including Snow Pea, Seashore Springtail; Entomobryid Springtail family including Ainsley's Springtail; Globular

Cockroaches

Blattid Cockroach Family including Oriental Cockroach (Asiatic Cockroach, Shad Cockroach), and American Cockroach (Waterbug); Blattellid Cockroaches family including German Cockroach (Croton Bug)

7.1383"

 $Rotting\text{-}wood\,termite\,family\,including\,Pacific\,Coast\,Termite;}\,Subterrane an$ termite family including subterranean termites. Not intended to replace structural treatments. For termites - Do not use this product as the sole source of control for active, structural infestations by subterranean termites; the purpose of this application is to kill workers or winged reproductive forms of termites which are present at the time of treatment It is not intended to provide structural pest control. It is not a substitute for mechanical alteration, soil and foundation treatment, but merely a supplement. For active, structural infestations by subterranean termites, this product can only be used to supplement a federally registered conventional product that is registered as a sole source for termite control. This product will not eliminate termite infestations or provide protection against future infestation.

Long-Horned Earwig family including Ring-legged Earwig, and Riparian Earwig; Common Earwig family including European Earwig

Grasshoppers & Crickets

Grasshopper Granily including Aztec Pygmy Grasshopper;
Short-Horned Grasshopper family including Creosote Bush Grasshopper,
Lubber Grasshopper, Painted Grasshopper, Carolina Locust, Dragon Lubber
Grasshopper (Gray Dragon), Three-banded Grasshopper (Banded Range
Grasshopper), Differential Grasshopper, Red-legged Locust, Spur-throated
Grasshopper, Two-stripped Grasshopper, Panther-spotted grasshopper,
Southeastern Lubber Grasshopper, Alutacea Bird Grasshopper, American Bird Grasshopper, Green Valley Grasshopper, Toothpick Grasshopper, Horse Lubber Grasshopper, Pallid-winged Grasshopper, and Great Crested Grasshopper (Dinosaur Grasshopper); Long-horned grasshoppers and Katydids family including Oblong-winged Katydid, Mormon Cricket, Short-legged Shield back Katydid, California Katydid, Angular-winged Stot (regged sine) adar kat yoli, Alaibrilla Arylid, Aliguda Winged Katydid, Keeled Shield-back Katydid, Nebraska Cone-head, Gladiator Katydid, Mescalero Shield-back Katydid, True Katydid (Northern Katydid), and Fork-tailed Bush Katydid; Camel Crickets family including Spotted Camel Cricket, Secret Cove Cricket, Jerusalem Cricket; True Crickets family including House Cricket (Cricket on the Hearth), Field Cricket, California Tree Cricket, Snowy Tree Cricket, and Black-horned Tree Cricket; Mole Cricket family including Northern Mole Cricket and European Mole Cricket

Plant Bug family including Adelphocoris Plant Bugs, Scarlet Plant bugs, Tarnished Plant Bugs, and Four-lined Plant Bug; Assassin bug family including Bee Assassins, Wheel Bug;
Ambush Bug family including Bee Assassins, Wheel Bug;
Ambush Bug family including Jagged Ambush Bug; Lace Bug
family including Oak Lace Bug; Seed Bug family including Small
Eastern Milkweed Bug; Long-necked Seed Bug, and Large Milkweed
Bug; Stilt Bug family including Stilt Bugs; Leaf-footed Bug family
including Florida Leaf-footed Bug, and Squash Bug; Scentless Plant Bug family including Eastern Boxelder Bug and Western Boxelder Bug; Stink Bug family including Green Stink Bug, Brown Marmorated Stink Bug Brochymenas, Two-spotted Stink Bug (Conspicuous Stink Bug), Harlequin Cabbage Bug (Calico Bug, Fire Bug), and Spined Soldier Bug

Cicada family including Periodical Cicadas, Dogday Harvestfly, and Grand Western Cicada

Treehoppers

Treehopper family, including Thorn-mimic Treehopper, Oak Treehopper, Buffalo Treehopper, and Locust Treehopper

Leafhopper family including Grape Leafhopper, Scarlet-and-green Leafhopper, Sharpshooter, Spotted Lanternfly*

Banded Thrips family including Banded-wing Thrips; Common Thrips family

Tiger Beetle Family including Beautiful Tiger Beetle, Dainty Tiger Beetle, Six-Spotted Green Tiger Beetle, and Dejean's Flightless Tiger Beetle; Ground Beetle family including Bombardier Beetle, Fiery Searcher (Caterpillar hunter), European Ground Beetle, Green Pubescent Ground Beetle, Common Black Ground Beetle, and Boat-backed Ground Beetle, Predacious Diving Beetle family including Small Flat Diving Beetle, Large Diving Beetle, and Marbled Diving Beetle; Whirligig Beetle family including Large Whirligig Beetle, and Small Whirligig Beetle; Hister Beetle family including Hister beetles; Rove Beetle family including Gold-and-brown Rove Beetle and Pictured Rove Beetle; Carrion Beetle family including Margined Burying Beetle, Tomentose Burying Beetle (Gold-necked Carrion Beetle), American Carrion Beetle, and Northern Carrion Beetle; Stag Beetle family including Elephant Stag Beetle, Agassiz's Flat-horned Stag Beetle, Reddish-brown Stag Beetle, and Rugose Stag Beetle; Bessbugs family including Patent-leather Beetle (Horned Passalus); Scarab Beetle family including European Chafer Beetle (adults), Flower Chafers (Fruit and Flower Chafers), Leaf Chafers, Rose Chafers, Tumblebugs, Goldsmith Beetle, Green Jumping Beetle, Brown Fruit Chafer, Glossy Pillbug, Grapevine Beetle, May Beetles (June Beetles), Ten-lined June Beetles, and Japanese Beetle; Metallic Wood-Boring Beetles including Golden Buprestid, Western Pine Borer (Sculptured Pine Borer), and Divergent Metallic Wood Borer (Flatheaded Cherry Tree Borer); Click Beetle family including Eastern Eyed Click Beetle Glier-eyed Click Beetlel, and Fire Beetles; Fireflies (Lightning Bugs) family including Pyralis Firefly, and Pennsylvania Firefly; Soldier Beetle family including Pennsylvania Leather-wing, and Downy Leather-wing, Net-winged Beetle family including Banded Net-wing, and Golden Net-wing; Spider Beetle family including Spider Beetles, and Texan Spider Beetle; Branch and Twig Borer family including Apple Twig Borer; Checkered Beetle family including California Checkered Beetle, Elegant Checkered Beetle, Slender Checkered Beetle, and Red-blue Checkered Beetle; Flat Bark Beetle family Checkered Beetle, and Red-Dide Checkered Beetle; Flat Bark Beetle family including Red Flat Bark Beetle; Pleasing Fungus Beetle including Rough Fungus Beetle; Ladybug Beetles including Multicolored Asian Lady Beetles, Two-spotted Ladybug Beetle, Nine-spotted Ladybug Beetle, Spottless "Nine-spotted" Ladybug, Convergent Ladybug Beetle, and Ash Gray Ladybug Beetle; Darkling Beetles including Horned Fungus Beetle, Broad-necked Darkling Beetle, Plicate Beetle, Yellow Mealworm Beetle, and Placehed Regular Fair Regular Beetle, Fire Regular Beetle, Plicate Beetle, Vellow Mealworm Beetle, and Ironclad Beetle; Fire-colored Beetles family including Fire Beetles; Blister Beetles family including Striped Blister Beetle, Arizona Blister Beetle,

.*Not Registered for Use by California.

Short-winged Blister Beetle, (Oil Beetle); Long-horned Beetle family including Black-horned Pine Borer, Yellow Douglass Fir Borer, Long-jawed Longhorn, Elder Borer (Cloaked Knotty-horn), Twig Pruners, Pine Sawyer (Spined-neck Longhorn), Golden-haired Flower Longhorn, Locust Borer, Black Pine Sawyer (White Spotted Sawyer), Cylindrical Hardwood Borer, Cottonwood Twig Borer, Cottonwood Borer, Giant Root Borers, Ribbed Pine Borer, California Laurel Borer (Banded Alder Borer), Ivory marked Beetle, Red Milkweed Beetle (Eastern Milkweed Longhorn), Notch-tipped Flower Longhorn, and Willow Borer; Seed Borer family including Bean Weevil; Leaf Beetle family including Dogwood Calligrapha, Milkweed Tortoise Beetle (Argus Tortoise Beetle), Dogbane Leaf Beetle, Milkweed Leaf Beetle, Willow Leaf Beetle, Spotted Asparagus Beetle, Clavate Tortoise Beetle, Spotted Cucumber Beetle, Waterlily Leaf Beetle, Swamp Milkweed Leaf Beetle, Three-lined Potato Beetle, Old-fashioned Potato Bug, Colorado Potato Beetle, and Girdled Leaf Beetle; Primitive Weevil family including Oak Timberworm Beetle (Primitive Weevil); Snout Beetles and Weevil family including Boll Weevil, Black Oak Acorn Weevil, Lesser Cloverleaf Weevil, Alfalfa Weevil, Pine Weevils, Rose Weevils, Agrave Billbug, and Stored-grain Billbug (Granary Weevil, Elephant Bug), Bark and Ambrosia Beetle family including Red Turpentine Beetle, Pine and Spruce Engraver Beetles, and Bark Beetles.

Bees, Ants, Wasps, Sawflies

Cimbicid Sawflies family including Rusty Willow Sawfly; Common Sawfly family including Northeastern sawfly; Horntail family including Pigeon Horntail, and Smokey Horntails; Stem Sawflies family including Raspberry Horntail; Braconids family including Braconid Wasps; Ichneumons family including Giant Ichneumons, Short-tailed Ichneumons and Red-tailed Including Grain Crimeunions, Short-Carlier Crimeunions and Red-Carlier Lichneumon; Torymid Wasp family including California Torymus; Chalcid family including Golden-yellow Chalcid; Gall Wasp family including Galifornia Oak Gall Wasps and Live Oak Gall Wasp; Pelecinid family including American Pelicini; Tiphiid Wasp family including Five-banded Tiphiid Wasp; Velvet-Ants including Thistledown Velvet-ant, Red Velvet-ant, and Cow Killer; Scoliid Wasp family including Scarab-hunter Wasp, and Digger Wasp; Ant family including Spine-waisted ants, Leafcutter ants, Texas Carpenter ants, Black Carpenter ants, Crater-nest ants, Crazy Ants, Tawny Crazy Ants, Texas Shed-builder ant (Acrobat Ant), Red Ant, Legionary Ant, Little Black IEXAS SNEO-DUILDER ANT (ACTOBAT ANT), RED ANT, LEGIDNARY ANT, LITTLE BIACK ANT, ANTA LANDS HONEY ANT, BIG-headed ANT, HONEY ANT, AND FIRE ANT, VESPID WASP family including Potter Wasp, Paper Wasps, Yellow Jackets, and Sandhills Hornet; Spider Wasps including Blue-black Spider Wasps, and Tarantula Hawk; Sphecid Wasp family including Thread-waisted Wasps, Eastern Sand Wasp, Steel-blue Cricket Hunter, Purplish-blue Cricket Hunter, Florida Hunting Wasp, Black and Yellow Mud Dauber, Cicada Killer, and Great Colden Dieger Wasps Reg Smiller including Plasters Reg. Vellow, faced Golden Digger Wasp; Bee families including Plasterer Bee, Yellow-faced Bees, Virescent Green Metallic Bees, Augochlora Green Metallic Bees, Alkali Bee, Mining Bee, Willow Mining Bee, Nevada Mining Bee, Clarkia Bee, Faithful Leafcutting Bee, Western Leafcutting Bee, Mason Bees, California Leafcutting Bee, Digger Bees, Western Cuckoo Bee, California Carpenter Bee

7.1383"

Crane fly family including Wood-boring Tupulid, Giant Western Crane Fly, and Crane Flies; Phantom Crane Flies including Phantom Crane Fly; Net-winged Midges including Comstock's Net-winged Midge; Phantom Midges; Black Flies including Black flies; March Flies family including March Flies; Fungus Gnats family including Fungus Gnats; Snipe Fly family including Gold-backed Snipe Fly; Stiletto Fly family including Stiletto Flies; Good-backed singerly, Stilettor Fly family including Stilettor Fless, Flower-loving Fly family including Flower-loving Fly; Mydas Fly family including Mydas fly; Robber Fly family including Bearded Robber Fly, Sacken's Bee Hunter, Giant Robber Fly, Bee Killer (Giant Robber Fly), and Robber Flies; Bee Flies including Bee Fly, Large Bee Flies, and Progressive bee Flies; Dance Flies including Dance Flies; Long-Legged Flies including Candel Stile Flower Flo Condylostylid Long-legged Flies, and Texan Long-legged Fly; Hover Fly family including Woollty Bear Hover Flies, Elongate Aphid Flies, Drone Flies Bulb Flies, American Hover Flies, and Toxomerus Hover Fly; Thick Headed Fly family including Thick headed fly; Pyrgotid Fly family including Pyrgotid

Flies; Fruit Fly family including Walnut Husk Fly, and Apple Maggot Fly; Seaweed Fly family including California Seaweed Fly; Marsh Fly family including Marsh flies; Shore fly family including Yellowstone Brine Fly; Pomace fly family including Yellowstone Brine Fly; Pomace fly family including Using Fly; Muscid Fly family including Using Fly; Muscid Fly family including House Fly, and Biting Stable Fly; Louse Fly family, Blow Fly family including Blue Bottle Fly, Screw-worm Fly, and Green Bottle Fly; Flesh Fly family including Flesh flies; Tachinid Fly family including Beelike Tachinid Fly, Early Tachinid Fly, Repetitive Tachinid Fly, Tachina Fly, and Fringe-legged Tachinid Fly

Folding trapdoor spider family including Turret spider; Tarantula family including Desert Tarantula; Trapdoor Spider family including California Trapdoor spider; Dictynid spider family including Branch-yip spiders; Spitting spider family including Spitting spiders; Violin Spider family including Desert Loxosceles, and Violin Spider; Comb-footed Spider family including American House Spider, and Sheet-web Weaver family including Hammock Spider; Orb Weaver family including Orb weavers, barn spider, Garden spider (cross spider), Marbled orb Weaver, Shamrock Spider, Six-spotted Orb-weaver, Silver Argiope, Black-and-Yellow Argiope, Crablike Spiny Orb Weaver, Bola Spider, Arrow-shaped Micrathena, and Golden-silk spider (calico spider); Large-jawed Orb Weaver family including Mabel Orchard Spider, Venusta Orchard Spider, Elongate Long-jawed Orb Weaver, and Long-jawed Orb Weaver; Funnel Web Weaver family including Grass Spiders; Nursery Web Spider family including Brownish-grey Fishing Spider, Spiters, Nulsery were Spiter raminy including brownisting by six Six-spotted Fishing spider, and Nursery Web Spider; Wolf Spider family including Burrowing Wolf Spiders, Carolina Wolf spider, Forest Wolf Spider, rabid wolf spider, and Thin-legged Wolf spider; Lynx Spider family including Jumping Lynx spider, and green lynx spider; Sac Spider family including Ant-mimic spiders; Wandering spider family including Wandering spiders; Giant Crab spider family including Huntsman spider, and Golden Huntsman spider; Selenopid Crab Spider family including Selenopid Crab Spider; Crab Spider family including Goldenrod spider (flower spider, red-spotted Crab Spider), Elegant Crab Spider, Thrice-banded Crab Spider; Philodromid family Spitely, Legain Crao Spited, The Charlest Jumping Spider, Finbutonina dia including Inconspicuous Crab Spiders; Jumping Spiders including Green Lyssomanes, Dimorphic Jumping Spider, Metaphid Jumping Spider, Ant-mimic Jumping spider, and Daring Jumping spider: Daddy-long-legs family including Eastern Daddy-long-legs and Brown Daddy-long-legs

Blacklegged tick, Brown dog tick, Deer tick, Lone Star tick

Psyllid, Whiteflies, Aphids, Mealybugs, Scale Insects
Psyllid family including American Alder Psyllid; Whitefly family including
Greenhouse Whitefly; Aphid family including Green Apple Aphid, Rosy Apple
Aphid, Rose, Pea and Potato Aphid, Root Aphid, Coudy-winged Cottonwood
Aphid, and Giant Willow Aphid; Woolly and Gall-Making Aphid family including Woolly Apple Aphid; Giant Scale Insects family including Cottony Cushion Scale; Armored Scale Insects family including Oyster Shell Scale; Mealybug family including Long-tailed mealybug; Cochineal Bug family including Cochineal bug.

Millipedes including soft millipedes, Florida millipede, Almond millipede, North American millipede and flat millipedes

