

QUICK KILL.

KILLS IN 24 HOURS

WORKS FOR 報 **3 MONTHS**







KILLS LISTED SURFACE & SUB-SURFACE INSECTS

TREATS UP TO 10,000 SQ.

NET WEIGHT **10 LB** (4.54 KG)

*Cis isomers 97% minimum, trans isomers 3% maximum

KEEP OUT OF REACH OF CHILDREN CAUTION SEE ADDITIONAL PRECAUTIONARY STATEMENTS ON BACK OF BAG


For outdoor use around residential sites only ‡Or your money back †Please refer to the insect lists found in the Directions for Use

KILLS IN 24 HOURS WORKS FOR ₩ 3 MONTHS

Distributed by:

CENTRAL

GARDEN & PET





1-800-265-0761 | gardendivision@central.com | amdro.com

EPA Reg. No. 59144-GA-1_A, 92162-MO-001_B EPA Est. No. 59144-GA-1_A, 92162-MO-001_B

design are trademarks of Central Garden & Pet Company. Scotts. Accugreen and Edgeguard are registered trademarks of OMS Investment: Inc. Speedygreen is a registered trademark of Everris International B.V.

treated area after application. as they appear. Thoroughly water the

or late evening hours. Avoid

disturbing ants. Treat new mounds

weather, 65 to 80 F, in early morning

mound. For best results apply in cool

granules to the surface of each apply 1 tablespoon or 3 teaspoons of

For **Fire Ant Mounds**: Uniformly

the treated area after application.

ft). For best results, thoroughly water

area of lawn or (4 lbs. per 1,000 sq.

Uniformly apply 1 lb. per 250 sq ft.

crickets, treat in late spring.

water the treated area after

Uniformly apply 1 to 2 lbs. per

of lawn. For best results, thoroughly

500 sq. ft. or 2 to 4 lbs. per 1,000 sq. ft

For best results, thoroughly water the

1,000 sq. ft. area of lawn using a

Uniformly apply 1 to 2 pounds over a

Application Rate

For best results in treating mole

when used as directed

use in measuring pesticides. NOTE: Food utensils such as tablespoons/teaspoons must not be used for food purposes after Repeat application if there are signs of renewed insect activity but not more than once every 7 days. For best results and to prevent further damage, treat the entire lawn, not just the browned areas.

Do not use any other weed, disease, or insect control products for at least one week after application. Water-in product after application. use a mechanical spreader. Spread this product uniformly over lawn using the suggested spreader settings. For best results, appear, or moths fly up from the grass when mowed).

SUMET

flower gardens, and vegetable gardens for the control of the insect pests listed in the table below. This product is for use on lawns, around home foundations (perimeter treatment), ornamental and GENERAL INSTRUCTIONS

Apply this product when insect activity or damage is first observed (lawn thins, brown spots

 Do not allow children or pets to enter the area until dry. product to enter any drain during or after application.

drainage to sewers, storm drains, water bodies, or aquatic habitat can occur. Do not allow the • Application is prohibited directly into sewers or drains, or to any area like a gutter where

Do not make applications during the rain.

Do not water to the point of run-off.

Apply this product directly to the lawn or garden area. Water treated area as directed on this label. It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

DIRECTIONS FOR USE the lawn or garden will help to prevent run off to water bodies or drainage systems.

Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. waters. Applying this product in calm weather when rain is not predicted for the next 24 hours not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface This pesticide is extremely toxic to fish and aquatic invertebrates. To protect the environment, do **ENVIRONMENTAL HAZARDS**

drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated eyes or clothing. Wash thoroughly with soap and water after handling and before eating, Harmful if absorbed through the skin. Causes moderate eye irritation. Avoid contact with skin,

> Hazards To Humans and Domestic Animals PRECAUTIONARY STATEMENTS

digestible fats and oils may increase absorption and so should be avoided. This product is a pyrethroid. If large amounts have been ingested, milk, cream or other

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-265-0761 for emergency medical treatment information. • Call a poison control center or doctor for treatment advice.

continue rinsing eye. Remove contact lenses, if present, after the first 5 minutes, then Hold eye open and rinse slowly and gently with water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

 Rinse skin immediately with plenty of water for 15-20 minutes. Take off contaminated clothing. **GIA T2817**



tin eyes

active ingredient into the grass and soil. Water-in product after application to release STEP 3



suggested spreader settings Spread uniformly over lawn using the STEP 2



from the grass when mowed). thins, brown spots appear, or moths fly up Observe insect activity or damage (ie. lawn

STEP 1 3 EASY STEPS

 $^{*}0.5$ lb/500 sq. ft. is equivalent to 1 lb/1000 sq. ft. there are signs of renewed insect activity. 2.0 lbs. for each 500 sq. ft. (2 ft. x 250 ft.). If necessary, repeat applications every 2 weeks only if

Home Foundation/Barrier Treatment

This product can be applied just before planting or after plants emerge.

activity, but not more than every two weeks.

Ornamentals & Flowers

may transmit Lyme Disease

Scorpions

SebeqilliM

Centipedes

Ground Pearl

Ground Beetle Heas (larvae and adults) Crane Flies (larvae)

Chiggers

Ticks (including Brown Dog Tick) and ticks that

Ants (including Argentine, Pharaoh (sugar), Imported Fire Ants, and Foraging Fire Ants)

Southern Mole Cricket, Short-Winged Mole

Mole Crickets (including Tawny Mole Cricket,

(czergeula linnuk gnibuloni) cliveeW

Annual Bluegrass Weevil (Hyperodes)

Big-Headed, and Allegheny mound)

Lawn, Cornfield, Ghost, Thief, Acrobat,

Ants (including Carpenter, Red Harvester, Red

Webworm, Tropical Sod Webworm)

Two-lined Spittle Bugs Striped Grassworm

Southwestern Corn Borer Springtails

Sowbugs

Spiders

Silverfish

Seed Maggots

Mealybugs

Lucerne Moth

Leaf Hoppers Mealybug Ladybird)

grasshoppers

European Crane Fly FSIMISS

Dichondra Flea Beetle Cutworm)

Wood Cricket)

Соскгоасћеѕ

Chinch Bugs Chiggers Bristletail

FIRE Brat Fiery Skipper

Webworms (including Sod Webworm, Grass

Twice-stabbed Ladybeetle, Asian Ladybeetle,

Ladybeetle, Thirteen-spotted Ladybeetle, Seven-spotted Ladybeetle, Two-spotted Ladybeeties (including Convergent Ladybeetle,

Cutworm, Granulate Cutworm, Variegated

Sand Field Cricket, Vocal Field Cricket, Southern

Western Striped Cricket, Eastern Striped Cricket,

Spring Field Cricket, Northern Wood Cricket

Crickets (including House Cricket, Common

Billbugs (including Bluegrass Billbug, Hunting

Armyworms (including Lawn Armyworm, Fall

Black Turfgrass Ataenius (adults)

Southeastern Field Cricket, Texas Field Cricket,

Cricket, Variable Field Cricket, Fall Field Cricket, Short-tailed Cricket, Arizona Cricket, Tropical

Cutworms (including Black Cutworm, Bronzed

White-footed, Crazy, Little Black, Small Honey, Pavement, Odorous, Pyramid, Black Turf,

Evenly apply a 2-foot-wide band of granules next to the house/home foundation. Use 0.5° to

Apply 0.5^* to 1 lb. evenly over each 500 sq. ft. area. Repeat if there are signs of renewed insect

renewed insect activity, but not sooner than two weeks after the first application • In New York State, do make a single repeat application of this product if there are signs of

100 feet of a water body (lake, pond, river, stream, wetland, or drainage ditch).

• In New York State, this product may NOT be applied to any grass or turf area (lawns) within

and use not in accordance with directions. Notice: To the extent consistent with applicable law, Buyer assumes all responsibility for safety product down any indoor or outdoor drain. $\textbf{partly filled:} \ \text{Call your local solid waste agency for disposal instructions.} \ \textbf{N} \text{ever place unused}$ reuse or refill this container. If empty: Place in trash or offer for recycling if available. If

PESTICIDE DISPOSAL AND CONTAINER HANDLING: Nonrefillable container. Do not **PESTICIDE STORAGE:** Store in a cool, dry, secure place. Do not store near seeds, fertilizers, Do not contaminate water, food or feed by storage or disposal. STORAGE AND DISPOSAL

Directions	Pounds of product per 1,000 sq. ft 1 lb. 2 lb. 4 lb.			2 bкездек
Shut off spreader anytime you stop. For complete coverage make each pass about five feet from the previous one, so some overlapping occurs.	b /I-9	Z/I-t	3	Scotts° Speedy Green° EdgeGuard° Rotary
Mow lawn first if grass is tall enough to bouch the boftom of the spreader. First apply two header strips. Operate spreader the longest way of the lawn. Overlap wheel marks to ensure uniform feeding. Shut off spreader when you shader the header when you have reached the header have reached the header have reached the header.	6	Ĺ	Z/I-S	Scotts® Accubreen® Drop Spreader

strips or stopped.

Spreader Settings Covers 5,000 sq. ft. Spreader Settings manufacturer's recommendations. Follow Application Directions in the table below. Calibrate the spreader settings based on the Spreading Instructions

4 to 6 inches of soil. Use 1.0 lb of product for a 500 sq. ft. area. Apply a single application before planting or after plants emerge and work into the top This product can be applied to the vegetables listed below.

Vegetable Gardens

GRANULES Lawn Insect GUICK KILL, DAGMA

crickets, grubs and cutworms spiders, ticks, chinch bugs, mole Not harmful to grasses¹ insects including ants, fleas, around your home Kills up to 100 listed

perimeter and flower gardens For use on lawns, ornamentals, home

Provides a barrier of protection