

GRANULES

KILLSTO 100 LISTED INSECTS

KILLS IN 24 HOURS

WORKS FOR 報 3 MONTHS









KILLS LISTED SURFACE & SUB-SURFACE INSECTS

TREATS UP TO 20,000 SQ.

NET WEIGHT **20 LB** (9.08 KG)

For outdoor use around residential sites only ‡Or your money back †Please refer to the insect lists found in the Directions for Use

KEEP OUT OF REACH OF CHILDREN CAUTION SEE ADDITIONAL PRECAUTIONARY STATEMENTS ON BACK OF BAG

ACTIVE INGREDIENT
*Bifenthrin OTHER INGREDIENTS 100.000%




KILLS IN 24 HOURS

WORKS FOR 報 **3 MONTHS**

1-800-265-0767 | gardendivision@central.com amdro.com

> EBY Reg. No. 59144-GA-1_A, 92162-MO-001_B Atlanta, GA 30339 **Distributed by:** Central Garden & Pet, Garden Division 1000 Parkwood Circle, Suite 700





CENTRAL GARDEN & PET

and use not in accordance with directions.

product down any indoor or outdoor drain.

Drop Spreader

Scotts® AccuGreen®

EdgeGuard® Rotary

Scotts® Speedy Green®

Spreader

Spreader Settings

Spreading Instructions

(Bell and Non-Bell)

and Spaghetti)

Cucurbit Vegetables

Sweet Corn

Tomatoes

Mustard, Eggplant, Head Lettuce, Peppers

Broccoli, Brussels Sprouts, Chinese Broccoli,

(Butternut, Calabaza, Hubbard including Acorn

Pumpkin, Summer squash (Crookneck, Scallop,

melon, Golden melon, Honeydew Melon, Pineapple

Chayote, Citron melon, Cucumber, Edible Gourd

Beans), Dry Beans (Black Eyed Peas, Cow Pea)

Snow Peas), Green Beans (Wax Beans, Snap

Green Peas, Edible Pods (Sugar Snap Peas,

Crop

Muskmelon (Cantaloupe, casaba, Crenshaw

Straightneck, Zucchini) Winter squash

melon, Santa Claus melon, Snake melon)

Chinese Cabbage, Kohlrabi, Chinese

Cabbage, Cauliflower, Cavalo Broccoli,

Do not contaminate water, food or feed by storage or disposal.

Notice: To the extent consistent with applicable law, Buyer assumes all responsibility for safety

partly filled: Call your local solid waste agency for disposal instructions. Never place unused

PESTICIDE STORAGE: Store in a cool, dry, secure place. Do not store near seeds, fertilizers,

STORAGE AND DISPOSAL

4-1/5

7 lp:

Pounds of product per 1,000 sq. ft

Spreader Settings Covers 5,000 sq. ft.

Follow Application Directions in the table below. Calibrate the spreader settings based on the

the header strips or stopped.

niform feeding. Shut off

Overlap wheel marks to ensure the longest way of the lawn.

header strips. Operate spreader the spreader. First apply two

enough to touch the bottom of

Mow lawn first if grass is tall

werlapping occurs.

orevious one, so some

about five feet from the

you stop. For complete

солегаде таке еасh pass

Shut off spreader anytime

Directions

Application

and wireworms

seed maggots, weevils

grasshoppers, grubs,

crickets, earwigs,

Ants, cutworms,

and wireworms

seed maggots, weevils

grasshoppers, grubs,

crickets, earwigs,

Ants, cutworms,

and wireworms

seed maggots, weevils

grasshoppers, grubs,

crickets, earwigs,

Ants, cutworms,

and wireworms

seed maggots, weevils

grasshoppers, grubs,

corn borer and weevils

эктумогт, grasshop-

bers, southwestern

corn borer, fall

earwigs, European

cutworms, crickets,

Ants, armyworms,

weevils and

till Harvest

Application

Days to Wail

seed maggots,

grasshoppers, grubs,

Pests

crickets, earwigs,

crickets, earwigs,

Ants, cutworms,

sbreader when you have reached

reuse or refill this container. If empty: Place in trash or offer for recycling if available. If

PESTICIDE DISPOSAL AND CONTAINER HANDLING: Nonrefillable container. Do not

 * 0.5 lb/500 sq. ft. is equivalent to 1 lb/1000 sq. ft. there are signs of renewed insect activity.

2.0 lbs. for each 500 sq. ft. (2 ft. x 250 ft.). If necessary, repeat applications every 2 weeks only if

Evenly apply a 2-foot-wide band of granules next to the house/home foundation. Use 0.5* to Home Foundation/Barrier Treatment

activity, but not more than every two weeks. Apply 0.5* to 1 lb. evenly over each 500 sq. ft. area. Repeat if there are signs of renewed insect This product can be applied just before planting or after plants emerge.

Ornamentals & Flowers

Ground Pearl

Ground Beetle

renewed insect activity, but not sooner than two weeks after the first application.

In New York State, do make a single repeat application of this product if there are signs of 100 feet of a water body (lake, pond, river, stream, wetland, or drainage ditch). In New York State, this product may NOT be applied to any grass or turf area (lawns) within

treated area after application. as they appear. Thoroughly water the disturbing ants. Treat new mounds or late evening hours. Avoid weather, 65 to 80 F, in early morning mound. For best results apply in cool granules to the surface of each apply 1 tablespoon or 3 teaspoons of For **Fire Ant Mounds**: Uniformly may transmit Lyme Disease Scorpions Ticks (including Brown Dog Tick) and ticks that

the treated area after application. ft). For best results, thoroughly water səbəqilliM Centipedes area of lawn or (4 lbs. per 1,000 sq. Imported Fire Ants, and Foraging Fire Ants) Uniformly apply 1 lb. per 250 sq ft. Ants (including Argentine, Pharaoh (sugar), Weevils (including Annual Bluegrass) Southern Mole Cricket, Short-Winged Mole Mole Crickets (including Tawny Mole Cricket,

Fleas (larvae and adults) application. Crane Flies (larvae) water the treated area after Annual Bluegrass Weevil (Hyperodes) of lawn. For best results, thoroughly Big-Headed, and Allegheny mound) Uniformly apply 1 to 2 lbs. per

crickets, treat in late spring.

For best results in treating mole

500 sq. ft. or 2 to 4 lbs. per 1,000 sq. ft Lawn, Cornfield, Ghost, Thief, Acrobat, White-footed, Crazy, Little Black, Small Honey, Pavement, Odorous, Pyramid, Black Turf, Ants (including Carpenter, Red Harvester, Red Webworm, Tropical Sod Webworm) Webworms (including Sod Webworm, Grass Sgud elthiq2 benil-owl Striped Grassworm Sowbugs Southwestern Corn Borer Springtails Spiders Silverfish Pillbugs Seed Maggots Mealybugs Lucerne Moth Leaf Hoppers Mealybug Ladybird) Twice-stabbed Ladybeetle, Asian Ladybeetle,

Ladybeetle, Thirteen-spotted Ladybeetle, Seven-spotted Ladybeetle, Two-spotted Ladybeetles (including Convergent Ladybeetle, Greenbug Grasshoppers Fruit Fly Fire Brat Fiery Skipper treated area after application. Earwigs

For best results, thoroughly water the Dichondra Flea Beetle

Black Turfgrass Ataenius (adults) Billbugs (including Bluegrass Billbug, Hunting

Cutworm) Cutworm, Granulate Cutworm, Variegated Cutworms (including Black Cutworm, Bronzed 1,000 sq. ft. area of lawn using a Uniformly apply 1 to 2 pounds over a Sand Field Cricket, Vocal Field Cricket, Southern Western Striped Cricket, Eastern Striped Cricket, Spring Field Cricket, Northern Wood Cricket Southeastern Field Cricket, Texas Field Cricket, Cricket, Variable Field Cricket, Fall Field Cricket, Short-tailed Cricket, Arizona Cricket, Tropical Crickets (including House Cricket, Common Chinch Bugs Chiggers Bristletail

Application Rate

use in measuring pesticides. NOTE: Food utensils such as tablespoons/teaspoons must not be used for food purposes after Repeat application if there are signs of renewed insect activity but not more than once every 7 days.

For best results and to prevent further damage, treat the entire lawn, not just the browned areas. Do not use any other weed, disease, or insect control products for at least one week after application. Water-in product after application. use a mechanical spreader. Spread this product uniformly over lawn using the suggested spreader settings. For best results, appear, or moths fly up from the grass when mowed). Apply this product when insect activity or damage is first observed (lawn thins, brown spots

SUMET

flower gardens, and vegetable gardens for the control of the insect pests listed in the table below. This product is for use on lawns, around home foundations (perimeter treatment), ornamental and GENERAL INSTRUCTIONS

product to enter any drain during or after application. drainage to sewers, storm drains, water bodies, or aquatic habitat can occur. Do not allow the • Application is prohibited directly into sewers or drains, or to any area like a gutter where Do not make applications during the rain.

Do not water to the point of run-off.

Do not allow children or pets to enter the area until dry.

USE RESTRICTIONS: Apply this product directly to the lawn or garden area. Water treated area as directed on this label.

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. DIRECTIONS FOR USE the lawn or garden will help to prevent run off to water bodies or drainage systems.

Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. waters. Applying this product in calm weather when rain is not predicted for the next 24 hours not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface

This pesticide is extremely toxic to fish and aquatic invertebrates. To protect the environment, do **EUVIROUMENTAL HAZARDS** clothing before reuse. drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated eyes or clothing. Wash thoroughly with soap and water after handling and before eating,

Harmful if absorbed through the skin. Causes moderate eye irritation. Avoid contact with skin,

Hazards To Humans and Domestic Animals PRECAUTIONARY STATEMENTS digestible fats and oils may increase absorption and so should be avoided.

This product is a pyrethroid. If large amounts have been ingested, milk, cream or other

for treatment. You may also contact 1-800-265-0761 for emergency medical treatment information. Have the product container or label with you when calling a poison control center or doctor, or going • Call a poison control center or doctor for treatment advice. Remove contact lenses, if present, after the first 5 minutes, then • Hold eye open and rinse slowly and gently with water for 15-20 minutes.

Call a poison control center or doctor for treatment advice.

• Rinse skin immediately with plenty of water for 15-20 minutes. Take off contaminated clothing. **GIA T2817**

active ingredient into the grass and soil. Water-in product after application to release

or clothing

It on skin

.0 suggested spreader settings Spread uniformly over lawn using the



STEP 2 from the grass when mowed). thins, brown spots appear, or moths fly up Observe insect activity or damage (ie. lawn STEP 1

3 EASY STEPS **NI NWAJ RUOY TABAT OT WOH**

4 to 6 inches of soil. Use 1.0 lb of product for a 500 sq. ft. area. Apply a single application before planting or after plants emerge and work into the top This product can be applied to the vegetables listed below. Vegetable Gardens

Lawn Insect GUICK KILL,

GRANULES

cutworms

mole crickets, grubs and spiders, ticks, chinch bugs, insects including ants, fleas, Kills up to 100 listed

• For use on lawns, ornamentals, home • Not harmful to grasses around your home Provides a barrier of protection

Armyworms (including Lawn Armyworm, Fall

J299

perimeter and flower gardens