



# GOPHER GASSER

## KILLS DESTRUCTIVE LAWN PESTS

POCKET GOPHERS, MOLES, GROUND SQUIRRELS,  
NORWAY RATS\*\*, SKUNKS, WOODCHUCKS\*

KEEP OUT OF REACH OF CHILDREN/MANTENER FUERA DEL ALCANCE DE LOS NIÑOS  
WARNING/ADVERTENCIA

\*Not for use on woodchucks in California.  
No lo use para marmotas en California.  
\*\*In North Carolina, this product can only be used on Norway rats.  
En Carolina del Norte, úselo para rata noruegas solamente.

ACTIVE INGREDIENTS:  
Potassium Nitrate ..... 45.0%  
Sulphur ..... 45.0%  
Carbon ..... 8.0%  
OTHER INGREDIENTS: ..... 2.0%  
TOTAL: ..... 100.0%

NET CONTENTS: SIX 0.75 OZ (21.27 g) GASSERS

TOTAL NET WT. 4.5 OZ (127.6 g)

CONTAINS

6 GOPHER GASSERS



KEEP OUT OF REACH  
OF CHILDREN  
WARNING

\*Not for use on woodchucks in California

\*\*In North Carolina, this product can only be used on Norway rats.

### PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS & DOMESTIC ANIMALS  
WARNING: Causes eye irritation. Harmful if absorbed through skin. Avoid contact with eyes, skin or clothing. Wash thoroughly with soap and water after handling. After ignition, cartridge produces toxic gases. Fumes may be harmful if inhaled.

### FIRST AID

If on skin or clothing:	- Take off contaminated clothing. - Rinse skin immediately with plenty of water for 15-20 minutes. - Call a poison control center or doctor immediately for treatment advice.
If inhaled:	- Move person to fresh air. - If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. - Call a poison control center or doctor immediately for further treatment advice.
If in eyes:	- Hold eye open and rinse slowly and gently with water for 15-20 minutes. - Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. - Call a poison control center or doctor immediately for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-265-0761 for emergency medical treatment information.

### ENVIRONMENTAL HAZARDS

This product is highly toxic to wildlife. Check all burrows for signs of non-target species. If present, do not treat burrows.

### ENDANGERED SPECIES CONSIDERATIONS

General Considerations  
It is a Federal offense to use any pesticide in a manner that results in the death of an endangered species. See the Endangered Species section in the Directions For Use section of this labeling. The killing of a member of an endangered species due to the use of gas cartridges may result in a fine and/or imprisonment under the Endangered Species act. All users of this product must determine whether endangered species are known to occur or might occur in the intended use areas. Consult the nearest U.S. Fish and Wildlife Service regional office or the appropriate State agency for current information on habitats occupied by endangered species.

### Specific Considerations

Use this product according to these instructions AND the instructions in the Endangered Species Protection Bulletin (see the "Use Restrictions" found under the "Directions For Use" section) to protect endangered and threatened species.

- Black-footed Ferret (*Mustela nigripes*): To limit risks to the black-footed ferret and to avoid destruction of its prey base, do not use this product within 4.3 miles (7 kilometers) of any prairie dog town unless the colony is an isolated black-tailed prairie dog town less than 80 acres in size or an isolated white-tailed or Gunnison prairie dog town less than 200 acres in size, or unless the town has been appropriately surveyed, using methods acceptable to the U.S. Fish and Wildlife Service, and found not to contain black-footed ferrets and found by the FWS not to be a suitable site for ferret reintroductions.
- Use of this product within the occupied habitats of the organisms listed immediately below is limited to qualified individuals who have been trained to distinguish dens and burrows of target species from those of nontarget species. Such individuals may be wildlife biologists, certified applicators, representatives of California state or Federal agencies, or agents of county agricultural offices or university extension offices.
  - Fresno kangaroo rat (*Dipodomys nitrosideus oileyi*) in Fresno, Merced, Kern, Kings, and Madera Counties, California.
  - Giant kangaroo rat (*D. ingens*) in Merced, Fresno, Kings, Tulare, Kern, San Luis Obispo, and Santa Barbara Counties, California.
  - Stephens kangaroo rat (*D. stephensi*) in Riverside, San Diego, and San Bernardino Counties, California.
  - Tipton kangaroo rat (*D. n. nitrosideus*) in Kings, Tulare, Kern, and Fresno Counties, California.
  - Point Arena mountain beaver (*Polydomes rube nigro*) in Mendocino County, California.
  - San Joaquin kit fox (*Vulpes macrotis macrotis*) unless authorized by the U.S. Fish and Wildlife Service in Alameda, Contra Costa, Fresno, Kern, Kings, Merced, Monterey, San Benito, San Joaquin, San Luis Obispo, Santa Barbara, Santa Clara, Stanislaus, or Tulare Counties, California.
  - Desert tortoise (*Gopherus agassizii*) in Arizona, southern California, southern Nevada, and southern Utah.
- Do not use this product within the occupied habitats of the endangered animals listed immediately below.
  - Morro Bay kangaroo rat (*D. heermanni morroensis*) in San Luis Obispo County, California.
  - Coachella Valley fringe-toed lizard (*Uma inornata*) in the Coachella Valley are of southern California.
  - Island night lizard (*Xantusia riversiana*) on Santa Barbara, San Clemente, and San Nicolas Islands, California.
- Blunt-nosed leopard lizard (*Gambelia silus*): Do not use this product within the occupied habitat of the blunt-nosed leopard lizard in Fresno, Kern, Kings, Merced, Monterey, San Benito, San Luis Obispo, Santa Barbara, Santa Clara, and Stanislaus Counties, California, from October 1 to April 15 unless a specific blunt-nosed lizard protection plan for this period is approved by the U.S. Fish and Wildlife Service and fully implemented. Use of this product in occupied habitat of this species from April 15 through September 30 is limited to daylight hours when air temperatures are 77-95°F (20-30°C).
- San Francisco garter snake (*Thamnophis sirtalis tetrataeneus*): Do not use this product within the occupied habitat of the San Francisco garter snake in San Mateo, San Francisco, Santa Clara, and Santa Cruz Counties, California from November 1 to March 30 unless a specific San Francisco garter snake protection program for this period is approved by the U.S. Fish and Wildlife Service and fully implemented. Use of this product in occupied habitat of this species under such an approved program or from April 1 through October 31 is limited to qualified individuals who have been trained to distinguish dens and burrows of target species from those of nontarget species. Such individuals may be wildlife biologists, certified applicators, representatives of California State or Federal agencies, or agents of county agricultural offices or university extension offices.
- Eastern Indigo Snake (*Drymarchon corais couperi*): Do not use this product within habitat type and locales known to support the Eastern indigo snake throughout Florida and the coastal plain of Georgia.

### CHEMICAL HAZARDS

Once ignited by the fuse, this cartridge will burn vigorously until completely spent and is capable of causing severe burns to exposed skin and clothes, and of igniting dry grass, leaves and other combustible materials.

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.  
USE RESTRICTIONS: FOR OUTDOOR USE ONLY. This product may only be used to control pocket gophers (*Thomomys*, *Geomys*, *Pappogomys*), moles (*Talpidae*), woodchucks (*Marmota*), Norway rats\*\* (*R. norvegicus*), skunks (*Mephitidae*), and ground squirrels (*Sciuridae*) in residential areas including lawns, ornamental and flower gardens, and open fields. Do not use cartridges inside or under buildings, homes, porches, or other structures due to possible fire hazard. Do not use cartridges where vegetation is dry and likely to catch fire. Do not use where use of gas cartridges is prohibited by local ordinances. AMDRO® Gopher Gasser will not control tree squirrels.

Endangered Species: It is a Federal offense to use any pesticide in a manner that results in the death of an endangered species. Use of this product may pose a hazard to endangered species or threatened species. When using this product, you must follow the measures contained in the Endangered Species Protection Bulletin for the area in which you are applying the product. To obtain Bulletins, no more than six months before using this product, consult <http://www.fws.gov/esa> OR CALL 1-844-447-3813. You must use the Bulletin valid for the month in which you will apply the product. Please note: Additional endangered or threatened species

ACTIVE INGREDIENTS:  
Potassium Nitrate ..... 45.0%  
Sulphur ..... 45.0%  
Carbon ..... 8.0%  
OTHER INGREDIENTS: ..... 2.0%  
TOTAL: ..... 100.0%

## KILLS POCKET GOPHERS, MOLES, GROUND SQUIRRELS, NORWAY RATS\*\*, SKUNKS, WOODCHUCKS\*

obligations are listed under "Endangered Species Considerations" on this label.  
When used as directed, the toxic gases produced will travel through unplugged burrow systems to kill the pests. In order for gas-producing cartridges to be effective on the pest species, a sufficient concentration of gas must be retained in the burrow system over a sufficient period of time. Damp soil tends to retain toxic gases within a burrow system better than dryer soil and, therefore, may enhance the effectiveness of this product. Use under conditions of extremely dry soil and vegetation can decrease product effectiveness and may increase the risk of starting brush fires.

### PRIOR TO APPLICATION

Carefully observe areas to be treated. Apply cartridges only to burrow systems or dens currently being used by target species. Do not apply cartridges to burrows or dens known or believed to contain non-target vertebrate animals. Ensure the cartridge will pass easily into opening (carefully clear tunnel entry, if necessary); and that enough soil, rocks, sod or other non-combustible material is on hand to close burrows.

For Moles (*Talpidae*): Identify the main runway for treatment, by looking for fresh conical mounds of dirt. Select the freshest of these, remove plug and dig in the direction of the main runway. If lateral to the mound is plugged, dig past the plug, or probe the ground around the plug until little or no resistance is found to locate the main runway.

For Norway Rats\*\* (*R. norvegicus*): Use only where burrows are isolated from homes, barns, and other man-made structures. DO NOT USE inside buildings or anywhere that gases could enter buildings.

For Pocket Gophers (*Thomomys*, *Geomys*, *Pappogomys*): Burrow systems usually consist of a main burrow 4-20" (10-50cm) below the ground surface and variable numbers of lateral burrows off of the main underground runway. Deeper branches off of the main runway are used as nests and for food caches. Burrow diameter ranges from 2-5" (5-12cm) and varies with the size of species. Locate characteristic fan-shaped soil mounds, which have signs of recent activity (e.g., recently excavated soil) at their plugged entrances. Remove soil plug from center of each burrow entrance. Return the next day to see if the burrow opening has been re-plugged, indicating active use by a pocket gopher. DO NOT treat burrows that remained unplugged. Find the horseshoe-shaped depression on one side of the plugged mound. Locate the main runway by probing the ground 15-18" (38-46cm) from the mound, starting on the same side as the depression. The main runway has been located when friction on the probe is released and it falls into the runway. Using a shovel or trowel, dig down to the main runway, taking care not to block it with soil. Treat all active mounds. One cartridge placed in the end of a runway should be sufficient. One or more additional cartridges must be used if treatment takes place in the middle of runway or if a runway is "T-shaped". Otherwise, there is a 50% chance that the pocket gopher will be on the untreated side of the plugged runway. Within 5 days of treatment, check area for appearance of fresh mounds, which could be the work of survivors or gophers that have moved into the treated area. Retreat all systems with evidence of new pocket gopher activity.

For Ground Squirrels (*Sciuridae*): Burrow design varies with the species, soil type, habitat, and climate. Burrows may be as much as 9' deep (nearly 3m). Burrow diameter varies with the body size of the species (and other factors) and may range from 2-10" (2-25cm). It is preferable to treat ground squirrel burrows during the spring breeding season. Observe the area in early morning before treating burrows to confirm that ground squirrels are actively using the site. DO NOT treat occupied burrows. DO NOT treat burrows with evidence that a snake or weasel has entered in pursuit of prey. Within 5 days of treatment, check area for appearance of live ground squirrels or opened burrow systems which could be the work of survivors or squirrels that have moved into the treated area. Retreat all systems with evidence of residual activity.


For Skunks (*Mephitidae*): Locate all entrances to dens. Do not treat dens under inhabited structures or buildings where toxic gases could reach humans, pets, or domestic animals including livestock.

For Woodchucks (*Marmota*): Locate all burrow entrances. Select one for use and close all other entrances with rock and dirt.

### APPLICATION

Important: Do not light the cartridge directly, or within 2 inches of face, nor hold a burning cartridge in your hand. The minimum fuse time is 5 seconds when it is lit at the tip. DO NOT INHALE TOXIC FUMES.

- Read and follow "Prior to Application" label directions for specific pests to be controlled.
  - Ensure the tunnel opening is large enough to easily fit cartridge (carefully clear tunnel entry, if necessary).
  - Place sufficient non-combustible material like soil, rocks or sod within reach so you can fully close treated burrow immediately after inserting cartridge.
  - Position yourself so that you can hold AMDRO® Gopher Gasser cartridge away from your body.
  - Remove tape strip and extend fuse out from cartridge.
  - Light tip of fuse with a match or lighter with a small flame.
  - Working quickly, insert the entire cartridge, fuse-end first, into one side of the open runway, burrow or den.
  - Immediately close the opening with non-combustible material. Tamp lightly to prevent gases from escaping. Immediately cover all openings from which gas escape is visible.
- For larger listed animals, or for longer or multiple runways, use more than one cartridge. Proceed to the next runway, burrow or den entrance and treat as above.



### STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.  
Pesticide Storage: Store only in original container, in a dry place inaccessible to children and pets.  
Pesticide Disposal & Container Handling: Nonrefillable container. Do not reuse or refill this container. If empty: Place in trash or offer for recycling if available. If partly filled: To dispose of unused cartridges, soak in water, crush, and bury in at least 6" of loose soil. Never place unused product down any indoor or outdoor drain.

\*Not for use on woodchucks in California

\*\*In North Carolina, this product can only be used on Norway rats

NOTICE: To the extent consistent with applicable law, seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. To the extent consistent with applicable law, buyer assumes all risk of use and/or handling of this product when such use and/or handling is contrary to label instructions.

Manufactured for:

AMBRANDS

1000 Parkwood Circle, Suite 700

Atlanta, GA 30339

EPA Reg. No. 73342-8

EPA Est. No. 088733-CHN-001

Made in China



Ampro, Central Garden & Pet, and Central Garden & Pet with design are trademarks of Central Garden & Pet Company.

© 2018-2024 Central Garden & Pet Company




1-800-265-0761 | gardendivision@central.com

9900041351AM1211V7-05/24



8 13576 00860 4





KILLS POCKET GOPHERS, GROUND SQUIRRELS,  
MOLES, SKUNKS, WOODCHUCKS & NORWAY RATS\*\*

## KEEP OUT OF REACH OF CHILDREN WARNING

ACTIVE INGREDIENTS:	
Potassium Nitrate	45.0%
Sulphur	45.0%
Carbon	8.0%
OTHER INGREDIENTS:	2.0%
TOTAL	100.0%

\*\*Not for use on woodchucks in California  
\*\*In North Carolina, this product can only be used on Norway rats.

## PRECAUTIONARY STATEMENTS

### HAZARDS TO HUMANS & DOMESTIC ANIMALS

**WARNING:** Causes eye irritation. Harmful if absorbed through skin. Avoid contact with eyes, skin or clothing. Wash thoroughly with soap and water after handling. After ignition, cartridge produces toxic gases. Fumes may be harmful if inhaled.

## FIRST AID

If on skin or clothing:	• Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor immediately for treatment advice.
If inhaled:	• Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. • Call a poison control center or doctor immediately for further treatment advice.
If in eyes:	• Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. • Call a poison control center or doctor immediately for treatment advice.
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-255-0761 for emergency medical treatment information.	

### ENVIRONMENTAL HAZARDS

This product is highly toxic to wildlife. Check all burrows for signs of non-target species. If present, do not treat burrows.

### ENDANGERED SPECIES CONSIDERATIONS

#### General Considerations

It is a Federal offense to use any pesticide in a manner that results in the death of an endangered species. See the Endangered Species section in the Directions For Use section of this labeling. The killing of a member of an endangered species due to the use of gas cartridges may result in a fine and/or imprisonment under the Endangered Species act. All users of this product must determine whether endangered species are known to occur or might occur in the intended use areas. Consult the nearest U.S. Fish and Wildlife Service regional office or the appropriate State agency for current information on habitats occupied by endangered species.

#### Specific Considerations

Use this product according to these instructions AND the instructions in the Endangered Species Protection Bulletin (see "The Use Restrictions" found under the "Directions For Use" section) to protect endangered and threatened species.

- Black-Footed Ferret (*Mustela nigripes*):** To limit risks to the black-footed ferret and to avoid destruction of its prey base, do not use this product within 4.3 miles (7 kilometers) of any prairie dog town unless the colony is an isolated black-tailed prairie dog town less than 80 acres in size or an isolated white-tailed or Gunnison prairie dog town less than 200 acres in size, or unless the town has been appropriately surveyed, using methods acceptable to the U.S. Fish and Wildlife Service, and found not to contain black-footed ferrets and found by the FWS not to be a suitable site for ferret reintroductions.
- Use of this product within the occupied habitats of the organisms listed immediately below is limited to qualified individuals who have been trained to distinguish dens and burrows of target species from those of nontarget species. Such individuals may be wildlife biologists, certified applicators, representatives of California state or Federal agencies, or agents of county agricultural offices or university extension offices.
  - Fresno kangaroo rat (*Dipodomys nitroideus exilis*):** In Fresno, Merced, Kern, Kings, and Madera Counties, California.
  - Giant kangaroo rat (*D. ingens*)** in Merced, Fresno, Kings, Tulare, Kern, San Luis Obispo, and Santa Barbara Counties, California.
  - Stephen's kangaroo rat (*D. stephensi*)** in Riverside, San Diego, and San Bernardino Counties, California.
  - Tipton kangaroo rat (*D. n. nitroideus*)** in Kings, Tulare, Kern, and Fresno Counties, California.
  - Point Arena mountain heaver (*Aphelandia rostr nigra*)** in Mendocino County, California.
  - San Joaquin kit fox (*Wulpes macrotis mitch*)** unless authorized by the U.S. Fish and Wildlife Service in Alameda, Contra Costa, Fresno, Kern, Kings, Merced, Monterey, San Benito, San Joaquin, San Luis Obispo, Santa Barbara, Santa Clara, Stanislaus, or Tulare Counties, California.
  - Desert tortoise (*Gopherus agassizii*)** in Arizona, southern California, southern Nevada, and southern Utah.
- Do not use this product within the occupied habitats of the endangered animals listed immediately below.
  - Morro Bay kangaroo rat (*D. heermanni morroensis*)** in San Luis Obispo County, California.
  - Coachella Valley fringe-toed lizard (*Uma inornata*)** in the Coachella Valley area of southern California.
  - Island night lizard (*Xantusia riversiana*)** in Santa Barbara, San Clemente, and San Nicolas Islands, California.
- Blunt-nosed leopard lizard (*Gambelia silus*):** Do not use this product within the occupied habitat of the blunt-nosed leopard lizard in Fresno, Kern, Kings, Merced, Monterey, San Benito, San Luis Obispo, Santa Barbara, Santa Clara, and Stanislaus Counties, California, from October 1 to April 15 unless a specific blunt-nosed lizard protection plan for this period is approved by the U.S. Fish and Wildlife Service and fully implemented. Use of this product in occupied habitat of this species under such an approved program or from April 1 through October 31 is limited to qualified individuals who have been trained to distinguish dens and burrows of target species from those of nontarget species. Such individuals may be wildlife biologists, certified applicators, representatives of California State or Federal agencies, or agents of county agricultural offices or university extension offices.
- Eastern Indigo Snake (*Drymarchon corais couperi*):** Do not use this product within habitat type and locales known to support the Eastern indigo snake throughout Florida and the coastal plan of Georgia.

### CHEMICAL HAZARDS

Once ignited by the fuse, this cartridge will burn vigorously until completely spent and is capable of causing severe burns to exposed skin and clothes, and of igniting dry grass, leaves and other combustible materials.

## DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.  
**USE RESTRICTIONS: FOR OUTDOOR USE ONLY.** This product may only be used to control pocket gophers (*Thomomys*; *Geomys*; *Pappogomys*), moles (*Talpidae*), woodchucks (*Marmota*), Norway rats\*\* (*R. norvegicus*), skunks (*Mephitidae*), and ground squirrels (*Sciuridae*) in residential areas including lawns, ornamental and flower gardens, and open fields. Do not use cartridges inside or over buildings, homes, porches, or other structures due to possible fire hazard. Do not use cartridges where vegetation is dry and likely to

catch fire. Do not use where use of gas cartridges is prohibited by local ordinances. **AMDRQ®** Gopher Gasser will not control tree squirrels. **Endangered Species:** It is a Federal offense to use any pesticide in a manner that results in the death of an endangered species. Use of this product may pose a hazard to endangered species or threatened species. When using this product, you must follow the measures contained in the Endangered Species Protection Bulletin for the area in which you are applying the product. To obtain Bulletins, no more than six months before using this product, consult <http://www.fws.gov/esp/bul> OR CALL 1-844-447-3813. You must use the Bulletin valid for the month in which you will apply the product. Please note: Additional endangered or threatened species obligations are listed under "Endangered Species Considerations" on this label.

When used as directed, the toxic gases produced will travel through unplugged burrow systems to kill the pests. In order for gas-producing cartridges to be effective on the pest species, a sufficient concentration of gas must be retained in the burrow system over a sufficient period of time. Damp soil tends to retain toxic gases within a burrow system better than dryer soil and, therefore, may enhance the effectiveness of this product. Use under conditions of extremely dry soil and vegetation can decrease product effectiveness and may increase the risk of starting brush fires.

#### PRIOR TO APPLICATION

Carefully observe areas to be treated. Apply cartridges only to burrow systems or dens currently being used by target species. Do not apply cartridges to burrows or dens known or believed to contain non-target vertebrate animals. Ensure the cartridge will pass easily into opening (carefully clear tunnel entry, if necessary), and that enough soil, rocks, sand or other non-combustible material is on hand to close burrows.

**For Moles (*Talpidae*):** Identify the main runway for treatment, by looking for fresh conical mounds of dirt. Select the freshest of these, remove plug and dig in the direction of the main runway. If lateral to the mound is plugged, dig past the plug, or probe the ground around the plug until little or no resistance is found to locate the main runway.

**For Norway Rats\*\* (*R. norvegicus*):** Use only where burrows are isolated from homes, barns, and other man-made structures. DO NOT USE inside buildings or anywhere that gases could enter buildings.

**For Pocket Gophers (*Thomomys*, *Geomys*, *Pappogomys*):** Burrow systems usually consist of a main burrow 4-20" (10-50cm) below the ground surface and variable numbers of lateral burrows off of the main underground runway. Deeper branches off of the main runway are used as nests and for food caches. Burrow diameter ranges from 2.5" (5-12cm) and varies with the size of species. Locate characteristic fan-shaped soil mounds, which have signs of recent activity (e.g., recently excavated soil) at their plugged entrances. Remove soil plug from center of each burrow entrance. Return the next day to see if the burrow opening has been re-plugged, indicating active use by a pocket gopher. DO NOT treat burrows that remained unplugged. Find the horseshoe-shaped depression on one side of the plugged mound. Locate the main runway by probing the ground 15-18" (38-46cm) from the mound, starting on the same side as the depression. The main runway has been located when friction on the probe is released and it falls into the runway. Using a shovel or trowel, dig down to the main runway, taking care not to block it with soil. Treat all active mounds. One cartridge placed in the end of a runway should be sufficient. One or more additional cartridges must be used if treatment takes place in the middle of runway or if a runway is "Y-shaped". Otherwise, there is a 50% chance that the pocket gopher will be on the untreated side of the plugged runway. Within 5 days of treatment, check area for appearance of fresh mounds, which could be the work of survivors or gophers that have moved into the treated area. Retreat all systems with evidence of new pocket gopher activity.

**For Ground Squirrels (*Sciuridae*):** Burrow design varies with the species, soil type, habitat, and climate. Burrows may be as much as 9' deep (nearly 3m). Burrow diameter varies with the body size of the species (and other factors) and may range from 2-10" (2-25cm). It is preferable to treat ground squirrel burrows during the spring breeding season. Observe the area in early morning before treating burrows to confirm that ground squirrels are actively using the site. DO NOT treat unoccupied burrows. DO NOT treat burrows with evidence that a snake or weasel has entered in pursuit of prey. Within 5 days of treatment, check area for appearance of live ground squirrels or opened burrow systems which could be the work of survivors or squirrels that have moved into the treated area. Retreat all systems with evidence of residual activity.

**For Skunks (*Mephitidae*):** Locate all entrances to dens. Do not treat dens under inhabited structures or buildings where toxic gases could reach humans, pets, or domestic animals including livestock.


**For Woodchucks (*Marmota*):** Locate all burrow entrances. Select one for use and close all other entrances with rock and dirt.

#### APPLICATION

**Important: Do not light the cartridge directly, or within 2 inches of fuse, nor hold a burning cartridge in your hand. The minimum fuse time is 5 seconds when it is lit at the tip. DO NOT INHALE TOXIC FUMES.**

- Read and follow "Prior to Application" label directions for specific pests to be controlled.
- Ensure the tunnel opening is large enough to easily fit cartridge (carefully clear tunnel entry, if necessary).
- Place sufficient non-combustible material like soil, rocks or sod within reach so you can fully close treated burrow immediately after inserting cartridge.
- Position yourself so that you can hold **AMDRQ®** Gopher Gasser cartridge away from your body.
- Remove tape strip and extend fuse out from cartridge.
- Ignite tip of fuse with a match or lighter with a small flame.
- Working quickly, insert the entire cartridge, fuse-end first, into one side of the open runway, burrow or den.
- Immediately close the opening with non-combustible material. Tamp lightly to prevent gases from escaping. Immediately cover all openings from which gas escape is visible.

**For larger listed animals, or for longer or multiple runways, use more than one cartridge. Proceed to the next runway, burrow or den entrance and treat as above.**



## STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.  
**Pesticide Storage:** Store only in original container, in a dry place inaccessible to children and pets.  
**Pesticide Disposal & Container Handling:** Nonrefillable container. Do not reuse or refill this container. If empty: Place in trash or offer for recycling if available. If partly filled: To dispose of unused cartridges, soak in water, crush, and bury in at least 6" of loose soil. Never place unused product down any drain or into outdoor drain.

\*Not for use on woodchucks in California

\*\*In North Carolina, this product can only be used on Norway rats

**NOTICE:** To the extent consistent with applicable law, seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. To the extent consistent with applicable law, buyer assumes all risk of use and/or handling of this product when such use and/or handling is contrary to label instructions.

Manufactured for:

**AMBRANDS**  
1000 Parkwood Circle, Suite 700  
Atlanta, GA 30339  
EPA Reg. No. 73342-8  
EPA Est. No. 088733-CIN-001  
Made in China

**AMDRQ.com**  
1-800-265-0761 | [gardendivision@central.com](mailto:gardendivision@central.com)  
990004135 | AM1211\_V7-05/24



Amdro, Central Garden & Pet, and Central Garden & Pet with design are trademarks of Central Garden & Pet Company.  
© 2018-2024 Central Garden & Pet Company



PARA ELIMINAR TALITZAS, ARDILLAS DE TIERRA, TOPOS,  
MORMOTAS Y RATOS NORUEGOS\*\*

## MANTENER FUERA DEL ALCANCE DE LOS NIÑOS ADVERTENCIA

ACTIVE INGREDIENTS:	
Potassium Nitrate	45.0%
Sulphur	45.0%
Carbon	8.0%
OTHER INGREDIENTS:	2.0%
TOTAL	100.0%

\*\*No lo use para marmotas en California.  
\*\*En Carolina del Norte, solo para rata noruegas solamente.

## AVISOS DE PRECAUCIÓN

### RIESGOS PARA HUMANOS Y ANIMALES DOMESTICOS

**AVISO:** Causa irritación en los ojos. Es dañino si se absorbe a través de la piel. Evite el contacto con ojos, piel o ropa. Lávese completamente con jabón y agua después de manipularlo. Después de ascenderlo, el cartucho produce gases tóxicos. Los vapores pueden ser tóxicos si se inhalan.

## PRIMEROS AUXILIOS

Si cae en la piel o ropa:	• Quite la ropa contaminada. • Enjuague la piel inmediatamente con suficiente agua durante 15 a 20 minutos. • Llame inmediatamente al centro de control de envenenamiento o al médico para obtener recomendaciones de tratamiento.
Si se inhala:	• Lleve a la persona al aire fresco. • Si la persona no respira llame al 911 o a una ambulancia, luego dé respiración artificial, preferiblemente de boca a boca, si es posible. • Llame inmediatamente al centro de control de envenenamiento o al médico para obtener recomendaciones adicionales de tratamiento.
Si entra en los ojos:	• Mantenga el ojo abierto y enjuague con agua después y con cuidado durante 15 a 20 minutos. • Si tienen fuertes de contacto, quíbelos después de lavarse con agua los primeros cinco minutos y después continúe lavando. • Llame inmediatamente al centro de control de envenenamiento o al médico para obtener recomendaciones de tratamiento.
Tenga a mano la etiqueta o el envase del producto cuando llame al centro de control de envenenamiento o al médico, o al buscar tratamiento. También puede comunicarse al 1-800-265-0761 para obtener información sobre tratamiento médico de emergencia.	

### RIESGOS AMBIENTALES

Este producto es altamente tóxico para los animales silvestres. Revise todas las madrigueras buscando indicios de especies que no son el objetivo. Si encuentra estos indicios, no aplique el tratamiento en las madrigueras.

#### CONSIDERACIONES PARA LAS ESPECIES EN PELIGRO DE EXTINCIÓN

##### Consideraciones generales

El uso de cualquier pesticida de manera que pueda ocasionar la muerte de una especie en peligro constituye un delito federal. Consulte la información sobre las especies en peligro en la sección "Indicaciones de uso" de esta etiqueta. La muerte de un miembro de una especie en peligro debido al uso de los cartuchos de gas se puede prevenir con mucha y/o prisión según la equidad la Ley de Especies en Peligro de Extinción. Todos los usuarios de este producto deben determinar si se conoce la existencia y posible existencia de especies en peligro en las áreas donde se aplicará el producto. Consulte en la oficina regional más cercana del Servicio de Pesca y Fauna Silvestre de Estados Unidos o la agencia estatal apropiada para conocer la información más actualizada sobre los habitats ocupados por especies en peligro.

##### Consideraciones específicas

Al usar este producto, siga estas instrucciones Y las instrucciones del Boletín para la protección de especies en peligro (véfere a las "Restricciones de uso" en la sección "Indicaciones de uso") para proteger las especies amenazadas y en peligro de extinción.

- Huías de patas negras (*Mustela nigripes*):** Para limitar los riesgos al huías de patas negras y evitar la destrucción de su área de caza, no use este producto a 4.3 millas (7 kilómetros) de cualquier colonia de perros de la pradera a menos que sea una colonia aislada de perros de la pradera de cola negra con un tamaño menor de 80 acres o una colonia aislada de perros de la pradera de cola blanca o Gunnison con un tamaño menor de 200 acres, o a menos que se haya investigado adecuadamente la colonia usando métodos aceptables al Servicio de Pesca y Vida Silvestre de EE.UU. (FWS) y que se haya determinado que no contiene hurones de patas negras y que el FWS no se un lugar adecuado para volver a introducir al huías.
- El uso de este producto en los habitats ocupados por los organismos detallados a continuación está limitado a personas calificadas que han sido capacitadas para distinguir entre guaridas y madrigueras de las especies objetivo con aquellas que no se desea eliminar. Esas personas pueden ser biólogos especializados en animales silvestres, aplicadores certificados, representantes de las agencias federales o del estado de California o agentes de oficinas agrícolas del condado o de extensión universitaria.
  - Rata canguro de Fresno (*Dipodomys nitroideus exilis*)** en los condados de Fresno, Merced, Kern, Kings y Madera, California.
  - Rata canguro gigante (*D. ingens*)** en los condados de Merced, Fresno, Kings, Tulare, Kern, San Luis Obispo y Santa Barbara, California.
  - Rata canguro de Stephen (*D. stephensi*)** en los condados de Riverside, San Diego y San Bernardino, California.
  - Rata canguro de Tipton (*D. n. nitroideus*)** en los condados de Kings, Tulare, Kern, Fresno, California.
  - Castor de montaña de Point Arena (*Ambrosia rostr nigra*)** en el condado de Mendocino, California.
  - Contra vuela de San Joaquin (*Vulpes macrotis mitch*)** a menos que esté autorizado el Servicio de Pesca y Vida Silvestre de EE.UU. en los condados de Alameda, Contra Costa, Fresno, Kern, Kings, Merced, Monterey, San Benito, San Luis Obispo, Santa Barbara, Santa Clara, Stanislaus, o Tulare, California.
  - Tortuga del desierto (*Gopherus agassizii*)** en Arizona, sur de California, sur de Nevada y sur de Utah.
- No use este producto en los habitats ocupados de los siguientes animales en peligro de extinción.
  - Rata canguro de Mono Bay (*D. heermanni morroensis*)** en el condado de San Luis Obispo, California.
  - Lagartija de dedos de Mono del Valle de Coachella (*Uma inornata*)** en el Valle de Coachella en el área del sur de California.
  - Lagartija Island night (*Xantusia riversiana*)** en Santa Barbara, San Clemente e Isla de San Nicolas, California.
  - Lagartija leopardo de nariz chata (*Gambelia silus*)** Este producto no se debe usar en el habitat ocupado de la lagartija leopardo de nariz chata en los condados de Fresno, Kern, Kings, Merced, Monterey, San Benito, San Luis Obispo, Santa Barbara, Santa Clara y Stanislaus, California del 1 de octubre al 15 de abril a menos que haya un plan de protección específico durante este período para la lagartija de nariz chata aprobado por el Servicio de Pesca y Vida Silvestre de EE.UU. y totalmente implementado. El uso de este producto del 15 de abril al 30 de septiembre en el habitat ocupado de esta especie está limitado a las horas con luz de día en que la temperatura del aire esté entre 77-95°F (28-30°C).
  - Serpiente jaretera de San Francisco (*Thamnophis sirtalis tetrastriatus*):** No se debe usar este producto en el habitat ocupado de la serpiente jaretera de San Francisco en los condados de San Mateo, San Francisco, Santa Clara y Santa Cruz, California del 1 de noviembre al 30 de marzo a menos que haya un programa de protección de la serpiente jaretera de San Francisco específico para este período aprobado por el Servicio de Pesca y Vida Silvestre de EE.UU. y totalmente implementado. El uso de este producto en un habitat ocupado de estas especies cuando están bajo tales programas aprobados a desde el 1 de abril hasta el 31 de octubre está limitado a personas calificadas que han sido capacitadas para distinguir entre guaridas y madrigueras de las especies objetivo con aquellas que no se desea eliminar. Esas personas pueden ser biólogos especializados en animales silvestres, aplicadores certificados, representantes de las agencias federales o del estado de California o agentes de oficinas agrícolas del condado o de extensión universitaria.
  - Serpiente indigo oriental (*Drymarchon corais couperi*):** No se debe usar este producto en los tipos de habitats y lugares conocidos donde se mantiene la serpiente indigo oriental a la larga de Florida y la planicie costera de Georgia.
  - Tortuga talpa (*Gopherus polyphemus*):** No se debe usar este producto en la cerca del habitat ocupado conocido de la tortuga talpa al oeste de los ríos Mobile y Tombigbee en Alabama.

### RIESGOS QUÍMICOS

Una vez encendida la mecha, este cartucho arderá fuertemente hasta agotarse y puede causar quemaduras graves en la piel expuesta y ropa, y encender la grama seca, hojas y otros materiales combustibles.

## INSTRUCCIONES

Es una violación a la ley federal usar este producto de forma inconsistente con su etiqueta.

**RESTRICCIONES DE USO:** PARA USO EN EXTERIORES SOLAMENTE. Este producto solo se puede usar para controlar las poblaciones de tozas o talitzas (*Thomomys*; *Geomys*; *Pappogomys*), topes (*Talpidae*), mormotas (*Marmota*), ratas pardas o ratas noruegas\*\* (*R. norvegicus*), mofetas o armillas (*Mephitidae*) y ardillas de tierra o marmotas (*Sciuridae*) en zonas residenciales incluyendo los espacios, jardines de flores ornamentales y campos abiertos. No use los cartuchos donde la vegetación está seca y propensa a incendiarse ni cerca de otras estructuras debido a los posibles riesgos de incendios. No use donde las leyes locales prohíben el uso de cartuchos de gas. El gas contra talitzas AMDRO® Gopher Gasser no es efectivo para controlar las ardillas que viven en los árboles.

Especies en peligro de extinción: El uso de cualquier pesticida de manera que pueda ocasionar la muerte de una especie en peligro constituye un delito federal. El uso de este producto puede representar un peligro para las especies amenazadas o en peligro de extinción. Al usar este producto, usted debe tomar las medidas mencionadas en el Boletín para la protección de especies en peligro para el área para la que se va a aplicar el producto. Para obtener los boletines, no más de seis meses antes de usar el producto, visite <http://www.fws.gov/esp/bul> OR CALL 1-844-447-3813. Usted debe usar el boletín válido para el mes en que va a aplicar el producto. Por favor, tenga presente la siguiente: En la sección "Consideraciones sobre las especies en peligro" de esta etiqueta se mencionan las obligaciones adicionales en cuanto a las especies amenazadas o en peligro.

#### ANTES DE LA APLICACIÓN

Observe cuidadosamente las áreas a tratar. Aplique los cartuchos únicamente a los sistemas de madrigueras o guaridas que usan las especies objetivo. No aplique los cartuchos en las madrigueras o guaridas que se sabe o cree que contienen animales vertebrados que no son el objetivo. Asegúrese de que el cartucho pasará fácilmente en la apertura (con cuidado la entrada del túnel claro, si es necesario), y que tiene a mano y que tiene a mano suficiente tierra, rocas, tepes u otros materiales no inflamables para cerrar las madrigueras.

**Para topes (*Talpidae*):** Identifique el paso principal para el tratamiento buscando montículos cónicos de tierra fresca. Seleccione el más fresco, quite el tapón y excave en dirección del paso principal. Si está tapado al lado del montículo, para localizar el paso principal excave después del tapón o explore el terreno alrededor del tapón hasta que haya poca o ninguna resistencia.

**Para ratas marmotas o noruegas\*\* (*R. norvegicus*):** Use solo donde las madrigueras están aisladas de los hogares, graneros y otras estructuras fabricadas por el hombre. NO LO USE adentro de edificios o en cualquier lugar donde los gases pueden entrar en los edificios.

**Para talitzas (*Thom occidentalis*, *Thom orientalis*, *Thom del sar*):** los sistemas de madrigueras consisten usualmente de una madriguera principal 4-20 pulgadas (10-50 cm) bajo la superficie de la tierra y cantidad variable de madrigueras laterales que salen del paso principal subterráneo. Se usan ramificaciones del paso principal más profundas como nidos y para almacenar alimentos. El diámetro de la madriguera es de 2 a 5 pulgadas (5-12 cm) y varía dependiendo del tamaño de la especie. Localice las características montículos de tierra en forma de abanico que tengan señales de actividad reciente (por ejemplo, tierra excavada recientemente) en las entradas tapadas. Quite el tapón de tierra del centro de cada entrada de madriguera. Regrese al día siguiente para ver si la abertura de la madriguera está tapada nuevamente. lo que indicará el uso activo de una talitza. NO trate las madrigueras que continúan sin tapar. Encuentre la depresión en forma de herradura que está en un lado del montículo tapado. Localice el paso principal explorando el suelo alrededor de 15-18 pulgadas (38-46 cm) del montículo, iniciando en el mismo lado que la depresión. Se localiza el paso principal cuando se libera la fricción en la vuelta de exploración y se en la pista. Use una pala o pala de jardinería, excave en el paso principal teniendo cuidado de no bloquearlo con tierra. Aplique tratamiento en todos los montículos. Debería ser suficiente colocar un cartucho al final de un paso. Si el tratamiento se realiza a la mitad de un paso o si el paso tiene forma de T, se debe usar uno o más cartuchos adicionales. De lo contrario, hay una probabilidad de 50% de que la talitza esté en el lado sin tratamiento del paso tapado. En los siguientes 5 días después del tratamiento, revise el área para buscar montículos frescos, que pueden haber hecho los sobrevivientes o talitzas que se trasladaron al área tratada. Vuelva a aplicar tratamiento en todos los sistemas que tengan evidencia de nueva actividad de talitza.

**Para ardillas terrestres (*Sciuridae*):** el diseño de la madriguera varía con la especie, tipo de tierra, habitat y clima. Las madrigueras pueden tener hasta 9 pies (casí 3 m) de profundidad. El diámetro de la madriguera varía con el tamaño del cuerpo de las especies (y otros factores) y puede estar entre 2 a 10 pulgadas (2-25 cm). Es preferible tratar las madrigueras de las ardillas terrestres durante la temporada de reproducción en primavera. Observe el área temprano en la mañana antes de tratar las madrigueras para confirmar que las ardillas terrestres están usando activamente el lugar. NO aplique tratamiento en madrigueras desocupadas. NO aplique tratamiento donde haya evidencia de que entró una cubeta o una cornucopia detrás de una presa. En los siguientes 5 días después del tratamiento, revise el área para buscar ardillas terrestres vivas o sistemas de madrigueras abiertas que pueden ser el trabajo de los sobrevivientes o ardillas que se trasladaron al área tratada. Vuelva a aplicar tratamiento en todos los sistemas que tengan evidencia de actividad residual.

**Para armillas o mofetas (*Mephitidae*):** localice todas las entradas a la guarida. No trate las guaridas que están bajo estructuras o edificios fabricados donde los gases tóxicos pueden llegar a los humanos, mascotas o animales domésticos, incluyendo ganado.


**Para marmotas (*Marmota*):** localice todas las entradas de la madriguera. Seleccione una y cierre todas las demás con piedras y tierra.

#### APLICACIÓN

**Importante: No encienda el cartucho directamente ni encienda la mecha a menos de 2 pulgadas (5,08 cm) del cartucho. No sostenga el cartucho ardiente en su mano. El tiempo mínimo de la mecha es de 5 segundos cuando se enciende en la punta. NO INHALE GASES TÓXICOS.**

- Lea y siga las instrucciones "Antes de la aplicación" de la etiqueta para ver las plagas específicas que desea controlar.
- Asegúrese de que la abertura del túnel sea lo suficientemente grande para insertar fácilmente el cartucho (después con cuidado la entrada del túnel si es necesaria).
- Coloque suficiente material no combustible como tierra, rocas o césped a su alcance para cerrar totalmente el túnel tratado inmediatamente después de insertar el cartucho.
- Colóquese en una posición que le permita sostener el cartucho del gasificador contra talitzas AMDRO® Gopher Gasser lejos de su cuerpo.
- Retire la tira de cinta y extienda la mecha para separarla del cartucho.
- Encienda la punta de la mecha con un fósforo o encendedor con llama pequeña.
- Rápidamente, inserte el cartucho completo (el extremo de la mecha primero) en un extremo del túnel, cavidad o madriguera abierta.
- Cierre inmediatamente la abertura con un material no combustible. Humedezca ligeramente para evitar el escape de los gases. Cubra inmediatamente todos las aberturas por donde vea que se escape el gas.

**Para animales más grandes, o para túneles más largos o múltiples, use más de un cartucho. Proceda hacia el próximo túnel, cavidad o madriguera y repita el tratamiento mencionado arriba.**



## ALMACENAMIENTO Y ELIMINACIÓN

No contamine el agua, los alimentos o los piensos al almacenar o eliminar.  
**Almacenamiento del pesticida:** Conservar solo en el envase original, en un lugar seco e inaccesible para los niños y las mascotas.  
**Pesticidas Disposición y Manejo de Contenedores:** Evase desechable. No vuelva a usar o llenar este envase. Si está vacío: Colóquelo en la basura o entréguelo para reciclar si está disponible. Si está parcialmente lleno: Para deshacerse de los cartuchos no utilizados, de inmersión en agua, aplástelo, y enterrar en por lo menos 6" de tierra suelta Nunca coloque el producto no utilizado en ningún drenaje interior o exterior.

\*\*No lo use para marmotas en California.

\*\*En Carolina del Norte, solo para rata noruegas solamente.

**AVISO:** En consonancia con la ley correspondiente, el vendedor no emite ninguna garantía explícita o implícita en cuanto al uso de este producto que no sea la indicada en la etiqueta. En consonancia con la ley correspondiente, el comprador asume todos los riesgos derivados del uso y/o la manipulación de este producto cuando dicho uso y/o manipulación no se corresponda con las instrucciones de la etiqueta.

Fabricado para:

**AMBRANDS**  
1000 Parkwood Circle, Suite 700  
Atlanta, GA 30339  
EPA Reg. No. 73342-8  
EPA Est. No. 088733-CIN-001  
Hecho en China

**AMDRQ.com**  
1-800-265-0761 | [gardendivision@central.com](mailto:gardendivision@central.com)  
990004135 | AM1211\_V7-05/24



Amdro, Central Garden & Pet, and Central Garden & Pet with design are trademarks of Central Garden & Pet Company.  
© 2018-2024 Central Garden & Pet Company

**AMDRO** **GOPHER GASSER**

**KEEP OUT OF REACH OF CHILDREN  
WARNING**

IMPORTANT: SEE PRECAUTIONARY STATEMENTS ON BACK OF PACKAGE AND FOLLOW ENTIRE DIRECTIONS FOR USE AND USE RESTRICTIONS ON ENCLOSED INSERT BEFORE USING.

AFTER IGNITION, CARTRIDGE PRODUCES TOXIC GASSES. FUMES MAY BE HARMFUL IF INHALED.

5 SECONDS MINIMUM FUSE BURN TIME WHEN LIT AT TIP

**MANTENER FUERA DEL ALCANCE DE LOS NIÑOS  
ADVERTENCIA**

IMPORTANTE: CONSULTE LOS AVISOS DE PRECAUCIÓN EN LA PARTE DE ATRÁS DEL EMPAQUE Y SIGA TODAS LAS INSTRUCCIONES Y RESTRICCIONES DE USO QUE ESTÁN EN EL FOLLETO ADJUNTO ANTES DE USAR EL PRODUCTO.

DESPUÉS DE ENCENDERLO, EL CARTUCHO PRODUCE GASES TÓXICOS. LOS VAPORES PUEDEN SER TÓXICOS SI SE INHALAN.

TIEMPO MÍNIMO DE 5 SEGUNDOS PARA QUE SE QUEME EL FUSIBLE CUANDO SE ENCIENDE EN LA PUNTA.

ACTIVE INGREDIENTS:	
Potassium Nitrate	45.0%
Sulfur	45.0%
Carbon	8.0%
OTHER INGREDIENTS:	2.0%
TOTAL:	100.0%

Manufactured for:  
AMBRANDS  
1000 Parkwood Circle, Suite 700  
Atlanta, GA 30339  
1-800-265-0761

Amdro is a registered trademark of Central Garden & Pet Company.  
990004135 | AM1211\_V7 - 05/24

EPA Reg. No. 73342-B  
EPA Est. No. 088733-CHN-001  
Made in China